







20 Grains make I Geruple) 3 Somples - 1 - 1 Dram 3
8 Drams - 1 - 10unce 3 12 Ounces _ 1 Pound to IS Semissis or half g.b. quantum libet as much as If quantum sufficit - vufficient John Elliot, The Medical Pocket Book, I England, 1791, with the autograph of Jere (1744/98), author of History of New Ha founder of Massachusetts Historical Soci work reprinted in America, rare Jeremy Belknap

Medical Pocket-Book.

CONTAINING

A short but plain Account of the SYMPTOMS, CAUSES, and METHODS of CURE, of the DISEASES incident to the HUMAN BODY.

Including fuch as require

SURGICAL TREATMENT:

Together with the

VIRTUES and Doses of Medicinal Compositions and Simples.

Extracted from the best Authors, and digested into Alphabetical Order.

GOOGOOGO TO TO THE SECOND COOM

THE TENTH EDITION,
WITH ADDITIONS AND CORRECTIONS.

By JOHN ELLIOT, M. D.

BOSTON:

rinted by JOSEPH BUMSTEAD, for DAVID WEST, No. 36, Marlborough-Street, and E. LARKIN, jun. No. 50, Cornhill. 1791.



AND THE STATE OF BYEAR ENDINGERS OF THE

SIR JOHN PRINGLE, BART,

PHYSICIAN TO THEIR MAJESTIES;

FELLOW OF THE ROYAL COLLEGES OF

PHYSICIANS AT LONDON AND

EDINBURGH;

AND OF THE

ROYAL SOCIETIES OF LONDON

AND PARIS, &c. &c.

THIS WORK

IS MOST RESPECTFULLY INSCRIBED,

BY HIS FAITHFUL

AND DEVOTED

HUMBLE SERVANT,

THE AUTHOR.

Great-Marlborough-Street, Aug. 12, 1781. FOR JOHN PHINGS, HARRY SERVES OF THE STREET AND A CONTROL AND SERVED AND A CONTROL AND SERVED AND S

MONAT COLUMNIA COLUMNAN COLUMNA COLUMNA

CINCARI CHV TOSMANYA SMI AN GRATICANI ARTARA SMI AN

JOHN MA and

Const Richmond Consts Light 22, 1923



PREFACE.

THE advantages of a publication of this kind to the practitioner in medicine, are ('tis prefumed) too obvious to need enumeration; I shall, therefore, only premise a few necessary observations relative to the work itself.

- I. The doses of medicines, &c. are fet down as for adults; and therefore, the prescriber will easily vary them for other ages. For example, children of two years old may, in general, take about a third part of the doses for adults, or a little more; and fo in proportion for other ages, the constitution being also considered.
- II. Where it could conveniently be done, I have mentioned the intervals at which the medicines are to be acpeated: but these may be varied at the preferiber's discretion. For brevity's sake I have in A 2 many

many places, used the word frequenter; by which I would have it understood, that every three, four, six, eight, &c. hours are meant, according to the exigency of the case.

III. When any particular indication is to be observed, the classes (as cathartics, diuretics, astringents, &c.) may be referred to; under which heads, as many of the compositions and simples as were judged necessary, are arranged.

IV. The limits affigned to the work would not admit of formulæ; nor do I think them by any means necessary. The doses of all the articles usually administered are set down; and I will not suppose, that there is any medical practitioner, who cannot give directions for making them into draughts, boluses, or other forms that may be required. The vehicles should be either persectly innocent, (as aq. pura, syr. simp. &c.) or of similar virtues with the medicines to be given. And if two articles of the same virtues (cathartics, for example)

are employed, it scarce need be mentioned, that the dose of each must be proportionably less. It may be added, that ingredients of a volatile nature should be given in a liquid form, lest their virtues exhale; very nauseous ones should, if convenient, be given in the form of pills or boluses.

V. In the alphabet I have, for the most part, made use of English names; or at least, have usually referred from those

where technical ones are employed.

This work was drawn up originally for my own use, and in the course of practice, I have sound it very convenient to refer to occasionally. It was the opinion of those medical Gentlemen who saw the M.S. that it would be equally useful to others, and this was my motive for making it public. If it assists the practitioner's memory in matters already known, (at least till he has leisure to consult larger works, where pecessary) my design is answered.



THE

Medical Pocket-Book.

ABORTION.

SYMPTOMS. Pain in the back, loins, and lower part of the belly; shiverings; a flux of blood from the womb; nausea, anxiety, palpitation of the heart, syncope, an opening and moisture of the ostincæ. Most commonly happens between the second and sourth months of pregnancy, though it may occur later. It may be occasioned by frights, sails, strong emetics or cathartics, or by any violent commotion of body or mind.

TREATMENT. Bleeding is necessary if the pulse will bear it. The medicines should be of the tonic and sedative kind, as tinct. rosar. or decoct. cort. with tinct. theb. The body should be kept lax with clysters, or otherwise; the diet light broths, and the like; and the patient should be kept very still and quiet in bed.

If, nevertheless, abortion follows, discontinue the corroborants, and give spermaceti medicines

with tinct, theb, &c, as after child-birth.

ABSCESS.

ABSCESS.

TREATMENT. Common abscesses may be eated by bringing them forward with proper traplasms, and when ripe, opening, digesting, id cicatrizing. Avoid evacuations during the ppuration, unless inflammation require. See so Boils, Phlegmon, and Ulcer.

ABSORBENTS.

(Of acidity) Creta, magnesia, crabs eyes, coral, d all the testaceous powders; dose \ni or \lnot or \lnot or \lnot veral times in a day. Also in like manner, salet. vel absinth. gr. v. sal ammon. vol. gr. v. luted.

ÆTHIOPS MINERAL.

Alterative, vermifuge; dose 3j. ad 3sf. bis die.

AGARIC.

Outwardly applied stops hæmorrhages; strewed a venereal warts, it sometimes drives them away.

AGUE.

SYMPTOMS. The fit begins with cold shivergs; a small quick pulse; pain in the back ad head; nausea. To these succeed great heat ad sever, which terminate in sweats. The urine uring the fit, pale, clear, and without sediment; at in the interval, turbid, with a copious sediment a reddish colour.

In the Quotidian Ague the fit returns once in a

In the Tertian, every other day.

In the Quartan, the intermission is of two

whole days.

Wards a gentle cathartic. If the intermissions are not regular, saline sebrifuges should be administered till that objection is removed; then give the bark, in substance zj. every two hours during the intermission, adding tinct. theb. or other astringent if it runs off by stool. If the stomach will not bear the powder, give it in decoction, insustant or the extract in pills. Pulv. sl. chamæma chalybs. rad. serp. virg. elix. vitr. acid, or t. theb may be added according to circumstances. The repetitions may be less frequent after the fit has been missed once or twice. Vitr. cærul. gr. sl. dissolved in zj. of proof spirit, and given before the fit, has succeeded in some desperate cases.

ALKALINES.

Sal. tart. sal absinth. sal vol. ammon. sal. c. c; vol. gr. v. ad pj. Sp. vol. am. sp. sal ammon. sp. C. C. gut. xv. ad zj. See also Absorbents. To be given occasionally to destroy acidities in the stomach, &c.

ALUM.

Astringent, gr. iv. ad Ass. Externally astringent and cooling.

ALOES.

Cathartic and anthelmintic off. ad 3fl. Emmenagogue gr. v. ad x. bis die. Externally applied, Vulnerary.

ALTERAT IVES

ALTERATIVES.

See the preparation of mercury and antimony.

ny medicine that works a cure gradually, by rrecting the patient's habit of body, may be lled an alterative.

ANASARCA. See Dropfy.

ANEURISM,

Or an enlargement of an artery, (known by its

ilfation.)

TREATMENT. Bleeding, and proper evacuaons; with low spare diet. Lead may be bound wn pretty tight to the part. If these fail, course must be had to the operation of tying it.

ANGINA GANGRENOSA, Or putrid fore throat.

Symptoms. Giddiness in the head, alternateeat and cold, and at length, after some hours, onstant great heat: sometimes vomiting or arging; pains in the head; eyes instanted and atery, as in the measses; anxiety, faintness, sore roat, with florid colour; or else, a broad regular spot, of a pale white colour, surrounded ith red. On the second or third day the ce, neck, breast, and hands swelled, and as if ysspelatous; an efflorescence of many small red imples sometimes appear on the arms and other arts. The throat sloughs, ulcerates, and is very painful. painful. Offensive taste, and delirium towards:

night.

TREATMENT. Avoid bleeding and evacuation, gentle sudorifics excepted; and purgings; should be checked. Give first a gentle emetic, then medicines with contraverva and conf. cardiac; or decoction of bark. Gargle, or rather syringe the throat with decoct. pect. cum rad. contray. to which, vinegar, myrrh, and honey of roses may be added. If the sloughs do not separate, touch them with a rag dipped in a mixture of mel. ægypt. zj. and zij. of the preceding gargle; or gargle with water acidulated with sp. of salt. To restore the patient give bark and elix. vitrioli, with proper strengthening diet.

ANISEEDS.

Carminative gr. iv. ad gr. xv. or in infusion ad ziv.

ANODYNES.

Opium crude or extr. gr. \(\frac{1}{4}\) ad gr. ij. tinc. theb. gut. x. ad xl. or more, fyr. e mecon zj. ad zff. elix. pareg. gr. xx. ad zjff. ther. andr. & conf. dam. \(\partial ff.\) ad zj. philon. Lond. \(\partial ff.\) ad zff.

ANTHELMINTICS. See Worms.

ST. ANTHONY'S FIRE. See Erifypelas.

ANTIMONY prepared.

Alterative gr. v. ad 3ff. bis die.

ANTISPASMODICS.

ANTISPASMODICS.

See Musk and its Julep. Opiates, and the nervous medicines.

APHTHE. See Thrush, and Fever Apthose,

APOPLEXY.

SYMPTOMS. A fudden privation of fensation, and voluntary motion. The face red, and bloated; the mouth commonly open; the pulse trong and quick, especially at first; respiration

trong, and attended with morting.

TREATMENT. Eleed freely, and apply cuping-glasses to the head, neck, or between the applies; strong clysters and cathartics; blisters applied to the back and legs, and mapisins to the feet. Medicines of the warm ervous kind, as castor, valerian, camphire, assumented, and volutile salts, should frequently be dministered.

Authors distinguish between a sanguineous and ituitous apoplexy: The first arising from a urgidness of the vessels of the brain; the last, om serous matter in the ventricles. In the tter case, emetics of the antimonial kind are rviceable, though often hurtful in the former; leeding should also be prescribed with caution in the latter. A soft, mild, and opening vegetable egimen is prescribed for patients subject to the anguineous, and a contrary one, with exercise, or such as are recovered from a pituitous poplexy.

B

APPETITE

APPETITE Canine.

TREATMENT. If an acid in the stomach be the cause, vomit, and give testaceous powders, magnefia, or alcalis. If worms, give anthelmintics. If it be natural, or proceeds from other causes, oils, fat meats, and broths, milk, and flour diet will be proper. Also opiates; smoaking tobacco is likewife of fervice.

APPETITE, Loss of.
TREATMENT. If the stomach be foul, give a gentle emetic of ipecacuanha; and if necessary, a gentle cathartic alfo. If it proceeds from relaxation, the bitters, chalybeates, or bark, with or without elix. vitriol. If acidity and wind abound, join fal absinth. with the bitters. Sena, rhubarb, tart. folubil. or other laxatives, may occasionally be employed if necessary.

ASCITES. See Dropfy. ASSAFCETIDA. See Gum.

ASTHMA.

SYMPTOMS. A difficulty of breathing, attended with wheezing; it returns at intervals; is preceded by difinclination to motion, loss of appetite, oppression, flatulency, and frequent eructations. At length, the cheeks become red, the eyes prominent, and there is fuch an anxiety and sense of suffocation, that the patient can only breathe in an erect posture, and can scarce speak or expectorate. If he happens to fleep he fnores much; at the height of the fit is defirous of cool free air; fweats about the neck and forehead; coughs up a little frothy matter with great difficulty; no fever; pulse extremely small and weak; urine pale and copious; as the fit abates, an expectoration of mucus; the urine becomes higher coloured, and deposits a copious sediment.

TREATMENT. During the fit bleed, if age or weakness do not forbid; blister between the shoulders, or at the pit of the stomach; the body should be opened with gentle cathartics; then give the pectorals and expectorants, joined, if

necessary, with antispalmodics and sedatives.

ASTRINGENTS.

Alum gr. iij. ad xij. pulv. e succin. c. gr. v. ad Dj tinct. rosar. Zj. ad ij. elix. vitrioli acid gut. x. ad xx. vitriol. alb. gr. ij. ad v. colcothar gr. iij. ad Ass. see also Corroborant. For the bowels in purgings, &c. ext. lign. camp, gr. v. ad Aj. alum, elect. e scord. Ass. ad zjss. rhab. torresact. gr. v. ad xv. Ripe acid fruits, as currants, sloes, &c. are also good.

ATROPHY.

SYMPTOMS. A nervous consumption, or wasting of the body; without fever, or other hectic symptoms.

TREATMENT. This must be varied according to the cause. If the appetite and digestion be bad, give a gentle puke, and afterwards the bitters

With

with steel. If scorphulous, the steel with bark. If worms be the cause, anthelmintics. If lues venerea, mercurials and decoct. farfaparil. If hysterics or hypochondriaca, join chalybeates with pil. gummos. or other medicines of this class. If profuse evacuation, as fluor albus, the bark, chalybeate waters, farfaparilla, and exercise. If atrended with scurvy, the scorbutic juices, vegetable acids, and bark. If afthmatic fymptoms appear, the squill preparations, or other pectorals and antispasmodics, and blisters. Diet, however, must affift medicine in the cure; which, where nothing forbids, should be of the foft, nutritive, and Arengthening kind, and easy of digestion; as milk, calves feet, and other jellies, &c. The country air is good.

A FTENUANTS.

Alcalis (which fee.) Chalybeates. See Flor. Mart. and Deobstruents, Mercurials, fee Calomel, &c.

BALSAM of Capivi.

Corroborant, diuretic, gut. x. ad xxx.

BALSAM of Guaiacum.

Gut. xx. ad zj. for rheumatic complaints, foulness of skin, gleets, &c.

BALSAM Lucatellus's.

Dj. ad zj. for dysenteries, or coughs.

BALSAM of Sulphur.

Gut. iv. ad Dj. for coughs and afthmatic complaints.

BALSAM of Peru.

Gut. iv. ad xv. for nervous complaints and head-aches, gonorrhœas, asthmas, &c.

Gr. v. ad pj. for coughs, &c.

BALSAM Traumatic.

Gut. x. ad xxx. for inward wounds, bruises, or coughs; externally for cuts.

BARK.

Febrifuge, corroborant, stomachic, antiseptic Dj. ad zj. several times in a day, decoction, or extract in proportion. The red bark is the species that ought to be preserved. For the revival of this excellent medicine, we are indebted to the authors of the London Medical Journal.

BARK of Pomegranates.

Astringent, stomachic, in decoction zj. ad zij, frequenter.

BARK of Lemons and Oranges. Stomachic, in infusion zj. ad zij. bis die.

BALAUSTINE Flowers.
Aftringent, Off. ad 3fl. or in decoction.

BEARING DOWN. See Procidentia, and Prolapfus.

B 2 BENJAMIN.

BENJAMIN.

Gr. iv. ad xv. as a pectoral and afthmatic.

BITES and STINGS of venomous animals.

TREATMENT. To those of small insects, as gnats, bugs, wasps, &c. apply spt. C. C. milk, oil, hony, or vinegar; to that of the viper, its own fat, spt. C. C. or ol. amygd. soment, poultice, and bleed, if inflammation require; and give milk inwardly, with theriaca, sal C. C. or other antidote. For the rattlesnake's bite, give a spoonful or two of juice of plantain or horehound, and apply a tobacco leaf steeped in rum to the part.

BLADDER inflamed.

SYMPTOMS. Acute burning pain, and tension in the part; with fever, frequent inclination to go to stool, and constant desire to make water. TREATMENT. As directed for kidneys infiamed.

BLEEDING external.

Apply the Applies mentioned under that word. Bleed at the arm, purge, and give cooling medicines if necessary. Keep the patient cool.

BLEEDING at the Nose.

TREAT as in Bleeding external; vinegar, or other flyptic, may also be snuffed up the nostrils, or cotton plugs, dipped in them put up. The patient in this case also, must be kept cool.

BLIGHT. See Erisypelas.

BLOOD,

Salt & water.

BLOOD, spitting of, or Hamoptoe.

SYMPTOMS. A cough and short respiration attends. If the blood coughed up appears florid, liquid, and frothy, especially if the patient be consumptive, it argues a rupture of some vessel in the lungs; is attended with pain if occasioned by a fall, or bruise.

TREATMENT. Bleeding, laxatives, and flypics; as tr. rofar. tr. flyptic, alum, bark, balf. raumat. pulv. e succin. c. nitro, and occasionally an opiate; the feet bathed in warm water; a slen-

ler cooling diet, and no exercise.

BLOOD, Vomiting of.

SYMPTOMS. There is no cough when the blood comes from the stomach; a sense of weight and oppression precedes the vomiting; the stomach s often distended; the blood of a darker colour than in an hæmoptoe.

TREATMENT. As in the preceding case. Small doses of vitr. virid. calcin. are efficacious in this, and also in a rupture of a vessel in the intestines, and a consequent discharge of blood by stool.

BOILS.

TREATMENT. They are to be ripened with proper poultices or cataplasms; opened, digested with basilicon, &c. and healed with cerat. epulot. or the like. If sungous sless appears, destroy it with red precipitate, or other proper escarotic. Alteratives and edulcorants, with corroborants, if eccellary, should be given internally.

BORAX.

BORAX.

Emmenagogue, diuretic, gr. v. or x. Externally to aphthæ.

BREASTS inflamed.

SYMPTOMS. A few days after delivery the breafts fometimes feel uneafy, or painful, and

fwell, the milk stagnating.

TREATMENT. Gentle purges and cooling diaphoretics. Apply diachylum plaisters, or empl. saponac, warm cloths, or the like, to the parts, and to the axillæ. If inflammation, soment, and poultice with bread and milk. If suppuration, continue to poultice, and let it break of itself; after which, digest with poultices or otherwise, and sicatrize.

If fever appears during the inflammation, treat

it as directed for fever (milk.)

If the breafts are merely hard, fomentations, poultices, and oil rubbed over them, are best.

BRUISES. See Contufions.

BUBO. See Venereal Disease and Abscess.

BURNS. See Scalds.

CACHEXY. See Dropfy.

TREATMENT. Sal diureticus, squills, or other diuretics; cathartics; then chalybeates, bark, &c. with proper exercise.

CÆLIAC PASSION.

SYMPTOMS. A purging, or discharge of ali-

ment indigested, and somewhat like chyle, with

wasting of the body.

TREATMENT. An emetic; stomachic, binding, and warm strengthening remedies; as bitters, spec. arom. philon. Lond. extr. lig. camp. opium, lect. e scord. creta, &c. with proper diet.

CALAMINE.

In collyriums for inflamed eyes; and in ointement as a cicatrizer.

CALAMUS.

Aromatic, gr. v. ad xv. CALOMEL. See Mercury.

CAMPHIRE.

Diaphoretic, diuretic, antispasmodic, gr. iij. ad

CANCER.

Symptoms. A round unequal tumour, of livid colour, furrounded with varicose vessels, and seated in the glandular parts of the body. ome cancers are fixed, others moveable; some ale, others red and inflamed; sometimes they emain harmless for years, at others they increase astily; ulcerate, and discharge a settid sanious thor, and soon prove mortal.

TREATMENT. Bleeding; in case of inflamnation, cooling laxatives; salt water, extract icutæ, magnesia, soap, or the like, in small doses salteratives. Externally, apply lint only if the umour breaks. Hemlock somentations have

been

been found serviceable, as hathalso the herb called clivers. Calmness of mind and proper regimen must be observed; but extirpation with the knife where it can be done, is the only certain remedy. For Cancer in the womb, see Wamb.

CANTHARIDES.

Analeptic, diuretic, gr. ff. ad gr. ij.

CARAWAYS.

Carminative, gr. iij. ad off. or in infusion.

CARDAMOMS.

Carminative, gr. iij. ad off.

CARDIALGIA. See Heart-burn.

CARMINATIVES.

Sem. cardam. anisi and carui, gr. iij. ad 38. Ol. carui, anisi, anethi, and juniperi gut. j. ad iij. Philonium 9ss. ad 3ss. Spec. arom. gr. iv. ad 3ss. Tinct. cardam. vel aromat. gut. x. ad 3j.

CARUNCLES. See Urine.

CARUS.

TREATMENT as in the apoplexy, of which this is only a flighter degree.

CASTOR.

Nervous, antispasmodic, emmenagogue, gr. iv. ad Dj.

CASTOR OIL.

Laxative 3ss. ad 3j. also nephritic and icteric. CATARRH.

CATARRH. See Cough.

CATHARTICS.

Jalap gr. xv. ad zff. Sal Glaub. zff. ad zjff. Aloes pff. ad pj. Pil. ex colocynth pj. ad pjj. Clect. e scam. zj. ad zij. Pulv. e scammon. c. pj. ad pij. Extract cath. gr. xv. ad zff. Coloynth pff. ad gr. xxv. Rhubarb pj. ad pij. Inf. enæ, zj. ad ziv. Tinct. sacræ, senæ, and rhuarb zj. ad zij. Gamboge, and scammony, pff. d pj. Ol. ricini zff. ad zj.

CEPHALICS.

Spt. lav. c. vel fpt. vol. arom. gut. x. ad xl. ether a few drops. Also externally to the temeles. See Nervous, and Head-ach.

CHALK.

Absorbent, cardialgic Dj. ad Zj.

CHALYBS prepared.

Deobstruent, tonic, gr. ij. ad off.

CHAMOMILE flowers.

Stomachic, corroborant, gr. v. ad Dj. or in a fusion.

CHICKEN-POX. See Pox. CHILBLAINS.

If they are not broke, bathe them with sp. vin. amph. sp. terebinth, urine, brine, melted salt utter, or rub them with snow. If they break, rest them with warm digestives, and afterwards with cerat epulotic.

CHILD-BIRTH.

CHILD-BIRTH.

If there be a profuse discourge of the lochia, give cooling medicines. If a puerperal, or milk fever, treat them as under those articles. delivery, and the after-birth, sperma-ceti, with gentle opiates; and if necellar,, caltor, or other uterine remedy.

CHINA Root.

Alterative, antivenereal, antiforbutic, in decostion like farfaparilla.

CHLOROSIS. See Menses.

CHOLERA MORBUS.

SYMPTOMS. Violent vomiting and loofeness, from bile abounding in the flornach and bowels; fharp pains, gripings, and flatulency; thirst, heat, anxiety, pulse quick and unequal; cold fiveats, and at length syncope, and coldness in the extremities. Chiefly attacks young persons, and in the fummer and autumn.

TREATMENT. Large and frequent draughts of chicken broth, to the quantity of a gallon or two; and clysters of the same should be continually injected. If the vomiting continues, give an infusion of out bread, toasted brown, in water; afterwards einnamon water, conf. card. elect. e fcord. or the like, with tinct. theb. An emetic may be proper, as also ripe fruit, &c. See Colic bilious.

CINNABAR of Antimony.

Alterative, vermifuge gr. iij. ad Bj.

CINNAMON.

CINNAMON.

Astringent, carminative, aromatic, gr. iij. ad psf. or in infusion.

COLCOTHAR.

Astringent, styptic, corroborant, gr. iij. ad gr. viij. Also externally to hæmorrhages.

COLD and COUGH.

SYMPTOMS well known. In phlegmatic habits, the cough is moist, and in the hypochon-

driac and scorbutic, dry.

TREATMENT. Gentle and regular warmth; bleeding, if nothing forbids; and especially if a fever, or pain in the breast, &c. arise. Oily, or spermaceti medicines, or other pectorals, with opiates. If costive, proper laxatives; and if the stomach is loaded with phlegm, an emetic; where greasy medicines disagree, give pil. de styrace, pulv. e trag. c. squills, mucilag. cydon. lozenges of liquorice, &c. Blisters are very efficacious, as are also gentle sudorifics repeatedly taken; with diluents and saline febrifuges, when the perspiration is obstructed, and sever appears.

COLIC, Bilious.

SYMPTOMS. An acute pain, and obstruction in the intestines; accompanied with a vomiting of yellow or greenish bile. A bitter taste in the mouth, with great heat. The pain sometimes fixed about the region of the navel, sometimes all over the abdomen, at other times shifting from

one

one part to another. A pulsation and cold are likewise selt in the belly. Urine little or none. Frequently hoarseness, with thirst and sever.

Most prevalent in summer.

TREATMENT. Bleed, and give an emetic, with a large quantity of inf. flor. ch. or the like, to work it off; then give cathartics (pills chiefly, or ol. ricini) with opium, and opening emollient clysters. If these fail, put the patient into the warm bath. If the vomiting continues, saline draughts, with tinct. theb. Pills of gr. sl. or gr. j. of crude opium are most powerful in removing pain and spasm; and enable the cathartics to take the better effect.

COLIC, Hysteric.

SYMPTOMS. This is a disease to which women of lax and gross habits, and of irritable dispositions, chiefly are subject. It begins with violent pain in the region of the stomach, much green and yellow matter is vomited, and great lowness of spirits succeed; the pain goes off in a day or two, and frequently returns again in a few weeks with equal violence. Sometimes 'tis attended with, jaundice, which in a few days goes off.

TREATMENT. Avoid bleeding and purging unless plethora or costiveness indicate; clear the stomach with warm water, chamomile tea, &c. then give a carminative antispasmodic, with opium; (as castor, conf. card. philon. conf. paulin.) which may be occasionally repeated. Recruit

the

the strength with bark, bitters, air, exercise, chalybeates, &c. This disorder in men is called the Hypochondriac Colic, and should be treated in like manner.

COLIC Inflammatory, or inflammation of the bowels.

SYMPTOMS. A vehement burning fixed pain is felt in the parts most affected, with heat of the whole body, quick pulse, loss of strength, anxiety and restlessness.

TREATMENT. Bleed largely, and repeat it if necessary, and the pulse will bear it; procure frools with fost mild cathartics, as oi. ricini, oil mixtures with manna and tart. folub. or the like; or if these will not stay, with pills of extr. cath. and opium; emollient clyfters should be frequently given, and foft emollient liquids drank; the bowels should be fomented, bladders of water, bags of falt or oats heated, the skins of animals just killed applied to the part; and the warm bath should be prescribed if these prove ineffectual. In case of violent vomiting, give the saline draughts in the act of effervescence; and the pain and spasm, if violent, may be abated with pills of crude opium; blifters may be applied to the part affected; the fumes of tobacco thrown up the fundament, in case all other cathartics fail; or give; quickfilver.

COLIC Nervous, called also the Devonshire-Golic, Colic of Poietiers, dry Gripes, and the dry Belly-ach. SYMPTOMS. SYMPTOMS. Begins with a fense of weight or pain at the pit of the stomach, attended with loss of appetite, yellowishness in the countenance, a slight nausea, and costiveness; a vorniting of acrid slime, and green bile succeed; the pain often descends to the navel, shooting from thence to each side, with great violence; the intestines seem drawn to the spine, with convulsive spasms; the pain continues without remission for several hours together; pulse as in health; no sever, but rather lowness; when at length the pain abates, an odd tingling is selt along the spina dors, which extending to the legs and arms, they

become weak and paralytic.

TREATMENT. Vomit several times with warm chamomile tea, then give opium in pills or otherwife, as also by clyster; the pain and tension being removed by these, give gentle cathartics, tuch as ol. ricini, infus.sen.with salts, and the like; laxative clysters may also be prescribed; bleed, if the pulse be high; stools being procured, an anodyne, carminative, and antispalmodic may be given, as camphire, valerian, castor, conf. card, or the like, with opium. If the pain returns, have recourse again to opium pills; if paralytic symptoms come on, embrocate the limbs and spine with lin. vol. cum camphor, or other warm liniment. If convulsions, give musk with opium. After the diforder, bark, with bitters and rhubarb, should be ordered; together with a light, but nutritive diet. Flannel may also be worn round the waist. Painters

Painters and glaziers, who are subject to this disorder on account of the lead used in their work, rely on burnt gin, which I have feen them take with furprizing fuccess.

COLIC Stone, or Niphritis.

Symptoms. The colic which fometimes ariseth from the stone, may be known from the other kinds by the following signs: There is a fixed pain in the kidney, which is propagated to the genitals; stools afford present relief in other colics, but not in this; after eating, the pain is increased in others, but lessened in this; in others, the urine is thicker in the beginning than afterwards, but in this, it is at first clear and thin, and afterwards lets fall gravel, and other sediment.
TREATMENT. Turpentine clysters are given.

with good effect; the pil. sapon. or other nephritic anodyne; care also being taken to keep the body open. But for more on this head, fee the

article STONE.

COLIC Wind.

SYMPTOMS. A wandering pain in the bowels, with rumblings, which abates on the expulsion of air; the pain is not increased by pressure; thirst not extraordinary; and the pulse but little disturbed.

TREATMENT. Bleed, if the patient can bear t; stools must be procured by saline or other eathartics, and emollient clysters; previous to which, an emetic also may be given if judged necessary; and somentations, or bladders of hot water may be applied to the belly; as may C 2 likewise

likewise anodyne, spirituous, and aromatic embrocations, as balf. anod. cum ol. menth. or the like. A passage being procured, carminatives joined with opium may be administered; as philon. Lond. or the like.

COLICA meconialis. See Meconium,

COLOCYNTH.

Cathartic, gr. x. ad Dj.

COMA. See Carus.

CONFECTION Alkermes. Cordial, Dj ad 3j.

CONFECTION Cordial. Carminative, cordial, off. ad oij.

CONFECTION Democrat. 'Alexipharmic, anodyne off. ad oij.

CONFECTION Paulins. Anodyne, nervous, Holland Holland

CONSUMPTION Nervous. See Atrophy. CONSUMPTION Pulmonary, or Phthisis.

parts of the body; dry cough; quick small pulse; moderate heat; lassitude; faintness; night sweats; slying pains and stitches; uneasiness about the diaphragm and breasts; the expectorated matter purulent; sometimes bloody and offensive, with white round lumps. If the symptoms are violent,

a spitting of blood soon follows, which is thin, florid, and frothy; but it afterwards becomes paler, and the discharge changes at length into pus. Ulcers, or tubercles in the lungs, are the cause.

TREATMENT. Small repeated bleedings; blisters kept open, and setons; gum ammon. myrrh, or balf. Peruv. with nitre, and elix. pareg. or pil. e styrac. at night; decoct. cort. may be given twice a day with good effect. While tubercles are forming, with fever and inflammation, avoid heating medicines, and give cooling febrifuges; if little or no fever, chalybeates are very efficacious; in spitting of blood, bleed and give tinct. rosar. tinct. stypic, or the like, and keep the body gently open. For sweats, give elix. vitri. acid. or if that disagrees, the pearl julep; very gentle vomits of ipecacuanha may be given twice a week, if nothing forbids, to promote expectoration; change of air and gentle exercise are good; the diet should be light, mild, and nutritive; as jellies, broths, milk, butter-milk, &c. Dr. Simmons's excellent treatife on this subject may be confulted.

CONTRAYERVA.

Alexipharmic gr. v. ad 5j.

CONTUSIONS.

TREATMENT. Bathe the part with vinegar in. fapon. lin. vol. fpt. mindereri, or arquebusade; and if necessary, apply a poultice of oatmeal and vinegar.

vinegar. If there be inflammation, bleed, and purge, use emollient fomentations and cataplasms. If suppuration appears, apply proper topical remedies to forward it, and treat as in abscesses.

CONVULSIONS.

Symptoms. Involuntarily contractions of spasses of the muscles, and consequent motions of the parts which they serve. Children are most subject to this complaint, and women of delicate constitutions.

TREATMENT. If a plethora be indicated, bleed; if otherwise, avoid it; blisters, laxatives, and emollient clysters, if nothing forbid; the nervous and antispasmodic medicines, as assaffasetida, castor, camphire, musk, volatile salts, with opium, if necessary. If the bowels are convulsed, opium may be added to the clysters also. The parts may be rubbed externally with bals. anod. and lin. vol. or other warm anodyne embrocation: free air should be allowed to the patient.

In children, this complaint often proceeds from an acid in the prima via; magnefia is then proper. Worms may also occasion it, anthelmintics are in this case to be given; blisters, and anodynes, given

with caution, are good.

CORAL pp. Absorbent 3j. ad 3j.

Anthelmintic off. ad 3ff.

CORNS.

Cover them with mucilage, or others plaister, feer bathing them in warm water, and paring em as much as convenient away. Large easy oes should be worn, and a little cotton may be id over them, to prevent their being rubbed pressed.

CORROBORANTS. See Bark, and its preparations; fee also Tonics.

COUGH, Hooping.

SYMPTOMS. It affects children; who by afon that the phlegm is difficult of expectoraon, strain violently to bring it up; with a hooping noise, till they are almost suffocated ad convulsed.

TREATMENT. If inflammatory fymptoms, or ver appear, bleed; give oxymel vomits, or ther small doses of ipecac. or tart. emet. to ing up the phlegm. Keep the body rather x; and prescribe tinct. sulig. or other antispasodic; gentle sedatives of syr. e mecon. are sicacious; blisters, gentle sudorifics, and a spare, in, but nourishing diet. Bark may be given, ith castor, to support the patient's strength. But lange of air, frequently repeated, sometimes bes wonders in these cases.

RABS CLAWS, Absorbent Dj. ad 3j. CREAM

CREAM OF TARTAR.

Aperient, cooling, 3ff. ad 3ff. or more; or made into whey, or infusion.

CUTANEOUS DISEASES. See the articles Itch, Scurvy, and Tetters.

CUTS.

Apply balf. traumatic, afterwards drefs with some mild digestive, and then cicatrize. If an artery is wounded, it must be taken up, or the spunge tent applied. See Wounds.

DANCE, St. Vitus's.

Symptoms. Convulsions of the legs, arms, and head; inarticulate speech, and lolling out of the tongue; drawing one leg after, like an idiot; with variety of odd and ridiculous gestures. Chiefly affects the youthful.

TREATMENT. Émetics, cathartics, pulv. rad. valer. in large quantities; bark, chalybeates, feabathing; electricity, millepedes, and æthiops mineral, have been of use; blisters and bleeding, if judged necessary. If worms are the cause, give anthelmintics.

DEAFNESS.

TREATMENT. If it proceeds from hardened wax, fyringe the ears, and drop in ol. amygd. If the wax be deficient, ol. amygd. with a little ol.rolifmar, or thelike. If cold be the cause, warmth; the remedies used against colds, and syringing with some warm soft liquid. If it be nervous,

ve the nervous medicines, bark, &c. and apply ifters. Electricity has been found useful. In flammation, bleed, purge, blister, foment, pultice, and prescribe the pediluvia. If there be cers, use injections, with tinct. myrrh, honey, c. If insects are in the ear, fill the meatus ith warm oil or water, or throw in the sumes tobacco.

DECOCTION, Pectoral. Ad libitum.

DECOCTION, White. Absorbent and astringent, ad libitum.

DELIVERY. See Child-birth.

DENTITION. See Teething.

DEOBSTRUENTS.

Pil. ecphractic off. ad oj. Sal martis gr. ij. vj. Calomel gr. ij. ad v. See also Menses, vine, Castor, Hellebore, Elix. Myrrh, &c.

DIABÆTES. See Urine.

DIAPHORETICS. See Sudorifics.

DIARRHŒA.

SYMPTOMS. A purging, without much fickis or pain; succeeded by loss of appetite; and metimes nausea, fever, with weak pulse, dry in, and thirst.

TREATMENT. If it proceeds from any thing fensive in the bowels, give rhubarb first, either powder or tineture; and afterwards astringents

and

and abforbents, as jul. e cret. elect. e fcord. extr. lig. camp. pulv. e bolo, &c. with or without opium as you fee necessary. Starch glysters, with opium, elect. e fcord. &c. and bleeding, if necessary, may be prescribed.

If weakness of the bowels and indigestion occasion the complaint, the bitters, bark, or chalybeates. If obstructed perspiration, gentle

fudorifics should be interposed.

Sometimes a purging is an effort of nature, to relieve the constitution from offending matters, or it is critical. You must be careful to distinguish in these causes, as checking the purging may be followed by a fever, or other bad consequences.

In diarrhœas, vegetable and acescent diet should be refrained from; decoct. alb. is best for common drink, and rice victuals for food. When purgings proceed from a putrescent cause, ripe fruits

and antiseptics are proper.

DIURETICS.

Sal diuretic pj. ad pij. Camphire gr. iij. ad pss. Ol. junip. gr. ij. ad v. Rad. scil. gr. iij. ad x. Spt. terebinth gut. x. ad xx. Spt. nitr. d. zss. ad zj. Mucilag. gum arab. ad libit. tinct. canthar. gr. x. ad xxx.

DROPSY.

SYMPTOMS. A collection of water or ferum in some part of the body. Before the disorder is perfectly formed it is called *Cacheny*; when the lymph

lymph is accumulated in the cellular membrane it is called Anafarca, or Leucophlegmatia; when there is a collection of water in the abdomen, it is termed Afcites; if in the brain, Hydrocephalus; if in the breast, Hydrops pectoris; if in the womb, Hydrops uteri; and if in the scrotum, Hydrocele.

In common dropsies the legs usually swell, and a dent or pit remains for some time after pressing the slesh with the singer; the appetite abates; the face either bloats or becomes thin and pale; little urine is made; thirst, slow sever, shortness of breath, lassitude and heaviness; these symptoms obtain especially in the analarca and ascites. In the hydrocephalus, convulsions, squinting, shunning the light, opening of the sutures, and vomiting of bile, commonly attend, especially if the disease be far advanced.

TREATMENT. In the Anafarca and Ascites, purges with jalap, scammony, calomel, gamboge, elaterium, or the like, twice or thride a week. Tinct. cupri. squills, tinct. cantharid. sal diurctic. or sal absent. with bitter insusion. Emetics and sudorifics, blisters and scarifications, prescribed with due caution, have often good effect, otherwise they may do harm. Bark may be given if the strength slags, or consumptive symptoms appear. Weak liquids of all kinds used to be avoided, but are now allowed freely with success, especially after the drastic purges. Crystals of tartar given in large quantity, have had good effect.

mended. In this complaint bread is generally ordered to be toasted, and slesh to be baked, roasted, or broiled. Smoaking is of great use, and mustard seeds taken whole have often been of service. If the disorder ariseth from the too copious use of weak liquids, or obstructed perspiration, sudorifies are much to be depended upon. If from drunkenness, riding on horseback, and the use of wine, or rather geneva diluted with water, in moderate quantity, will be useful. If from a consumptive tendency, diuretics, joined with corroborants. If after great loss of blood, or from tedious severs, cathartics should not be used too freely, but chalybeate bitters should be chiefly relied on. In the Ascites, tapping must be performed, if the methods above recommended fail.

In Dropfy of the breast, diuretics are chiefly to be employed, cathartics only occasionally. Blisters applied below the breast, and even on the thighs or legs, and kept open, are also very effectual.

In the Hydrocephalus intermus, a salivation excited by mercury has of late been recommended, but its efficacy is not yet clearly ascertained; purges may be given, and medicines of the carminative and antispasmodic kind. Blisters, setons, and issues should also be prescribed.

In the Dropfy of the womb, emetics, stimulating clysters,

clysters, and occasionally, cathartics; diuretic, attenuant, and resolvent medicines should be given; and fomentations and vapour baths advised.

In the Hydrocele, let out the water with the trocar at the bottom part of the scrotum; but when water is contained in the cellular membrane, treat it as an Anafarca.

> DRY GRIPES, or Dry-belly-ach. See COLIC Nervous. DYSENTERY, or Bloody-flux.

SYMPTOMS. A discharge of mueus, blood, purulent matter by stool; violent gripings; pain in the loins, and anus; tenesmus, and fever.

TREATMENT. A rhubarb, or other gentle purge, and after proper evacuation, julep e cret. with spermaceti, and an opiate. Bleed, if you find it necessary; and give starch clysters with opium and aftringents; boluses of bees-wax and permaceti, or balf. locat. with el. e scord, are efficacious. Small doses (a grain or two) of pecac. have often succeeded; as hath also vitrum intim. cerat. in dofes of gr. v. with some opiate, occasionally repeated. Sudorifics are sometimes of use, by promoting a determination to the urface of the body. Emollient somentations, nd balfamic clysters with opium, will best emove the tenefmus.

The diet as in the diarrhoea. Milk with mut-ETT OODAYALIZAD

ton fuet boiled in it, and the fat afterwards taken off, is esteemed excellent.

DYSURIA. See Urine.

EAR-ACH. See Deafness.

EARTH, Japan. Astringent, Oss. ad 3st.

ELATERIUM. Cathartic in dropfies gr. j. ad iij.

ELECTUARY of Bayberries, 3ff. in clyfters.

ELECTUARY Lenitive; or of Cassia. Laxative zij. and zss.

ELECTUARY of Seammony. Cathartic, antirheumatic zj. ad zij.

ELECTUARY of Scordium.

Astringent 3ss. ad 3jss. also sudorific.

ELIXIR of Aloes.
Deobstruent, opening, vermifuge gut. xv. ad 3j.

ELIXIR of Myrrh compound. Emmenagogue gut. v. and 3ss.

ELIXIR Paregoric.
Asthmatic, anodyne gut. xx. ad zjss.

ELIXIR of vitriol acid.

Corroborant gut. vj. ad xx.

EMMENAGOGUES.

EMMENAGOGUES. See Deobstruents.

EMETICS.

Pulv. ipecac. əff. ad əff. Vin. ipec. zj. ad zij. Tart. emet. gr. j. ad iij. Vin. antim. zija ad zff. Vitriol. alb. əff. ad əj.

EMPYEMA.

SYMPTOMS. An enlargement of the cavity of the thorax, and cedematous foulness of the skin, and slesh of one side thereof; dry cough, and difficult breathing: It ariseth from matter formed, and lying loose in the thorax.

TREATMENT, may be much like that prefcribed in the Vomica. Medicines which promote absorption, and blifters may also be useful. But letting out the matter, where it can conveniently be done, is the best cure.

EMULSION, Common.

Diuretic, emollient, ad libitum.

EPILEPSY.

Symptoms. Languid pulse, pale countenance, and afterwards great pain in the head, with stupor and drowsiness, sometimes precede the sit; though it often comes on without these previous symptoms. The patient salls down suddenly, gnashes the teeth, froths at the mouth, uses many disagreeable gesticulations and distortions, and sometimes discharges involuntarily by stool and wine.

D 2

TREATMENT.

TREATMENT. If the vessels are full, bleed; emetics and laxatives are proper; warm, nervous, and attenuating remedies, as castor, valerian, assaciated, cinnabar, viscus querc. ol. animal. salc. c. &c. should be directed. Blisters kept open, and setons are very serviceable; slowers of zinc have succeeded well, a grain or two at a dose once or twice a day. To restore the strength, and prevent returns, the bark, or other corroborant, and the cold bath, with proper diet, air, and exercise.

ERYSIPELAS.

SYMPTOMS. The usual precursory symptoms of a fever; the face, or other parts affected, inflamed; with scurf, pimples, and blisters; heat, redness, itching and smarting; drowsiness and

difficulty of breathing commonly attend.

TREATMENT. Bleed, and give proper lenitives, and cooling diaphoretic febrifuges. Blifters, and cordial sudorifics, as camphire, contraverva, &c. if low pulse and malignancy require. The parts may be bathed with softening somentations, milk, &c.

ESCHAROTICS.

Merc. corrosiv. ruber, vitriol. cærul. the caustics, alum ust. verdigris, &c.

ETHIOPS. See Æthiops.

EXTRACT of Bark.

Corroborant, &c. gr. v. ad Dj.

EXTRACT

EXTRACT Cathartic. Gr. xv. ad pij.

EXTRACT of Chamomile, or of Gentian. Stomachic, gr. v. ad Hs.

EXTRACT of Guaiacum.

off. ad 3ff. Alterative, antirheumatic, anti-

EXTRACT of black Hellebore. Deobstruent, opening, gr. v. ad H.

EXTRACT of Hemlock.
Deobstruent, alterative, &c. gr. v. ad Dj.

EXTRACT of Jalap. Cathartic, Hist. ad His.

EXTRACT of Logwood.

Aftringent, gr. v. ad xv.

EXTRACT of Saturn. Externally, cooling, aftringent, &c. diluted.

EXTRACT of Savine. Emmenagogue gr. v. ad 9ff.

EXTRACT Thebaic. Anodyne gr. ff. ad gr. ij.

EYES inflamed, or Ophthalmia. Inflammation of the outward coats of the eye, tended with pricking pain, heat, pulfation, redefs, fwelling, and scalding tears.

TREATMENT.

TREATMENT. Bleed, purge, apply leeches to the temples, and blifter the back, or behind the ears; cooling febrifuges may also be given; direct collyriums, with vitr. alb. sacc. saturn. or alum; or if astringents disagree, warm milk, decoction of mallow leaves in milk or water, or other emollient somentation; or poultice of milk and bread, and bathe the feet in warm water. If it ariseth from mere weakness of the vessels, astringent collyriums, or alum curd; and bark, or other tonics, may be given internally. If Nervous, join valerian, castor, or the like, with the bark. If Scorbutic, or other humours attend, perpetual blisters or setons, with mercurial, or other proper alteratives. If Tubercles in the eye are the cause, anoint them with ung. cærul. and give small doses of calomel till they disperse.

For watery Eyes, gentle cathartics and alteratives, and wash them with brandy and water. For specks in the eye, blow lap. cal. sugar, of specks, tutty, white vitriol with sugar, or the like, through a small tube into the eye; or drop in

folutions of white vitriol, facc. faturn. &c.

FALLING SICKNESS. See Epilepsy, FEBRIFUGES.

Nitre gr. v. ad Dj. Saline draughts. Vin. antimon. gr. v. ad xx. Tart. emet. gr. \(\frac{1}{4}\) ad \(\frac{1}{2}\). Bark in the intermissions, zss. ad zj. tertia q. hora, decoction, or extract in proportion. See Sudorifics, &c.

FENNEL.

FENNEL.

Carminative, diuretic gr. iij. ad gr. xv. or in infusion.

FEVER Apthose, or the Thrush Fever in adults. SYMPTOMS. Fever, with ulcerations or aphthæ; distinguishable from the putrid fore throat by the whiteness of the sloughs, by the edges not being red, and by there being no shining redness over the fauces.

TREATMENT. Febrifuges, bark, and antiseptics; blitters if necessary. Proper laxatives, and detergent gargles. Vitrum antimon. cerat. has been found uteful. Borax, alum, or spt. vitriol, when mixed in small quantity with honey, are good to touch the aphtaæ with. See also Thrush.

FEVER Erysipelatous. See Erysipelas.

FEIRED FOUL MAN

SYMPTOMS. Shivering, followed by heat, and quick pulse; nausea, anxiety, restlessness, white and dry tongue, thirst, and sometimes pain in the head and back.

TREATMENT. If there be fulness of vessels, bleed; give proper laxatives, and if the stornach be foul, an emetic; saline febrifuges, with small doses of vin. antim. or tart. emetic. and if diaphoresis be wanted, add contrayerva. If delirious, blister, and give julep e camph. If colliquative sweats, the decost cort. with or without elix vitriol. If spasm and catching of the tendons, with low pulse, blister the arms and legs, and apply stimulating

stimulating plaisters to the feet, first bathing them with warm water. If hiccups, jul. e mosch. While the pulse is high, and inflammatory symp-

While the pulse is high, and inflammatory symptoms continue, the antiphlogistic treatment, cooling febrisuges, acids, and gentle aperients are proper. But care must be taken not to bring the patient too low, else he will want strength to carry him through the disease. Gentle opiates may be ordered at night occasionally in case of restlessness. If the pulse sinks, he should be supported with proper cordials. Costiveness should be avoided, and even a purging should be checked with caution, being generally either critical, or at least salutary. The present and past symptoms compared, will best direct the treatment in severs. Diluting acidulated liquids are proper; the sood may be panada, gruels, puddings, and chicken broth, with bread, but no stess. Food or liquids particularly longed for should be given. Care should be taken to avoid too great heat and impure air.

FEVER Intermittent. See Ague.

FEVER Miliary.

SYMPTOMS. Shivering, heat, lowness of spirits, oppression about the præcordia, sighing. On the third or fourth day the eruption (preceded by profuse sweat of a sourish smell, with tingling or pricking sensation) appears, chiesly on the neck, breast, and back; of little bladders resembling millet-seeds, (from whence the sever is denominated)

nated)

ated) which are either white or red, according the colour of the liquid they contain, and somemes both forts. The eruption being out, the imptoms usually abate; the urine, which before as pale, becomes higher coloured. The erupons commonly dry in about feven days, and the

TREATMENT. Bleed, if nothing forbids; out this should be done with great caution) and ve faline febrifuges, accompanied with laxatives d diaphoretics, as the symptoms may require. bilious matter be suspected in the stomach, an netic may be proper. If delirium appears, ifter; if low nervous symptoms, give proper rdials; if putrid ones and petechiæ, the hark; aphthæ, gargles with tinct. myrrh, tinct. rosar. el, or the like. Acidulated liquids, fruit, &c. ould be given in this disease, the air kept pure, d the room not too hot.

FEVER Milk.

It ariseth about the third or fourth day after livery, with swelling of the breasts, and pain poting towards the axillæ; the breafts fomehes are hard, hot, and inflamed; generally conues a day or two, then terminates in copious eats and discharge of urine.

TREATMENT. Bleed, if the inflammation be eat; keep the body open, and let the breasts be quently drawn. In case of hardness or inflamtion, emollient fomentations and poultices.

FEVERS

FEVERS mixed,

Are those wherein the symptoms of inflammatory, nervous, putrid, and other severs, are more or less blended together; and must therefore be treated according to their appearances.

FEVER puerperal.

SYMPTOMS. Begins in two or three days after delivery, with the usual febrile symptoms; and ariseth from an inflammation of the omentum and intestines; pain in the fore part of the head, slushing in the face, short breath, suppression of the lochia.

TREATMENT. In case of costiveness, emollient glysters; and if necessary, gentle cathartics; then mild diaphoretics, as saline draughts with vinantim. and proper diluents. Emetics exhibited early, have been known to have been attended with remarkable success in this disease in the Hotel Dieu at Paris.

FEVER, putrid or malignant.

Symptoms. A giddiness precedes the attack; then burning heat, sudden loss of strength, heaviness, lowness of spirits, watching, pulse weak, low, and unequal; anxiety, oppression at the præcordia, nausea, vomiting, noise in the ears, desirium, coma, catching of the tendons, a black dry tongue, and thin crude urine. Petechiæ often appear on the fourth, fifth, and seventh days.

TREATMENT. Avoid bleeding, unless the pulse indicates its necessity; give small doses of

tart.

parts,

tart. emet. or pulv. jacobin with julep e camp. and if necessary, rhubarb, or other gentle laxative should be occasionally interposed. An emetic in the beginning is often efficacious. Antiseptics, the bark, and cordials are proper; as are also blisters. Opiates should be cautiously administered, and purgings, unless critical, should be checked. Sinapisms to the feet are proper in case of stupor; antispasmodics and jul. e mosch. in case of convulsive symptoms, or hiccups. The liquids drank should be acidulated; fruit may be allowed, as may also wine; the air in the room should be kept as pure as possible, and not too warm; vinegar should be sprinkled on the floor. Picking the bed-clothes in this disease is a bad symptom.

FEVER remittent, (Simple.)

SYMPTOMS. The remittent fever differs from the continual and intermittent, in that after a certain number of hours it remits, or abates, but does not go off. It is also called the autumnal remitting Fever, the bilious Fever, the marsh

Fever, and the camp Fever.

Usually comes on suddenly, with debility, low-ness of spirits, chilliness, and other febrile symptoms; the hands tremble; countenance pale or yellowish; skin dry, breathing difficult, and pulse small and quick; great nausea succeeds. Vomiting of bile, and sometimes discharge of it by stool; tongue becomes soul, delirium sollows, moisture on the sace, and at length on the other

parts, when the remission follows. As the disorder increases the remissions are less, and at length scarcely perceptible; the mouth, teeth, and lips, covered with a black crust; tongue very dry and stiff, so that the patient's voice can hardly be heard.

TREATMENT. Bleed not but with great precaution; an emetic is sometimes proper; the body should be opened if necessary; then give saline febrifuges, to bring on regular intermissions; this done, prescribe the bark copiously, as

directed for the ague.

FEVER, scarlet. (Simple.)

SYMPTOMS. The usual precursory symptoms of a fever, viz. Shivering, heat, &c. then an efflorescence of a scarlet colour appears all over the skin, but does not rise above the surface; with heat, dryness, and itching. In three or four days it disappears, and the cuticle comes off in branny scales.

TREATMENT. Cooling faline febrifuges, with gentle laxatives and diaphoretics, if necessary.

FEVER, fcarlet, (Malignant.)
Symptoms. Chillines, languor, sickness, oppression, succeeded by heat, nausea, vomiting, fore throat, quick pulse, and difficult breathing; the tonfils inflamed and ulcerated. On the third day the efflorescence appears.

TREATMENT. Saline febrifuges, joined occasionally with contrayerva, or other gentle diaphoretic; blifters on the back and

bleed

bleed and purge with great caution. If costiveness, clysters are best; and if inflammatory symptoms, cupping or leeches; if a putrid tendency appears, the bark, with rad. serp. virg. or rattlesnake root; tinct. theb. or tinct. cinnam.may be added if it purges; gargles of tinct. myrrh, tinct. rosar. mel, spt. salis, or the like, which should be injected with a syringe.

FEVER, sow, or nervous.

Symptoms. Shiverings, laffitude, wearinefs, fighing, pale desponding looks, great anxiety, depression of spirits, pain and giddiness of the head, white tongue (sometimes red) with a yellow or brownish list running along the middle of it, but no thirst; nausea, difficulty of breathing, pulse weak, quick, and unequal; urine limpid and pale; pain and coldness in the back part of the head; drowsiness. These symptoms are all worse towards night; if a delirium, not violent, but a muttering to themselves; sometimes miliary eruptions and sweats appear, but seldom afford relief.

TREATMENT. Gentle cordial diaphoretics; previous to which, an emetic, if nothing forbids. Costiveness should be removed bygentle laxatives; bleeding should be refrained from, unless urgent symptoms indicate; blisters; if an intermission appears, the bark; if convulsions, musk with castor or other antispasmodics; if aphthæ, gargles, with tinct. myrrh, honey, tinct. rosar. decoct. cort. alum, or the like; and in case of much

phlegm,

phlegm, oxymel scillit. or ipecac. to bring it up. Care should be taken after the sever, to restore the patient by proper nutritious diet, chalybeate waters, bark, &c.

FEVER, Worm. See Worms.

FEVERFEW.

(In posset drink) sudorific.

FLOODING. See Menses and Blood.

FLOWERS Balaustine. Astringent, gr. v. ad xx.

FLOWERS of Benjamin.

Pectoral, asthmatic, gr. ij. ad vj.

FLOWERS of Rosemary and Lavender.

Cephalic, in infusion zij. ad ziv.

FLOWERS Rose.

Astringent, gr. v. ad pj. or in infusion.

FLOWERS of Steel.

Deobstruent, corroborant, attenuating, gr. ij. ad viij.

FLOWERS of Sulphur.

Alterative, pectoral, opening, 9j. ad 3ij.

FLOWERS of Zinc.

Antiepileptic gr. \(\frac{1}{2}\) ad ij. emetic; outwardly, cooling and drying.

FLUOR

GANGRENE.

Symptoms. In some cases, when a part is violently inflamed, the inflammation and pain suddenly disappear; the integuments turn pale, blue, livid, or blackish; the skin becomes flaccid and pitting, with bladders filled with yellow or reddish ichor; and this is termed Gangrene.

reddish ichor; and this is termed Gangrene.

TREATMENT. Scarify and foment the parts with fotus comm. and spt. vin. camph. apply poultices of oatmeal and beer, and give cordial medicines, with plenty of bark. In gangrene of the toes, particularly in old people, opium has

been found particularly serviceable.

GARLICK.

Expectorant, diuretic, emmenagogue, a chive or two at a dose, or in infusion.

GENTIAN.

Stomachic, see Infusion (bitter.)

GINGER.

Cordial, aromatic, carminative, gr. v. or in infusion.

GLEET.

See Venereal Disease; but if it does not proceed from that cause, the same treatment will be proper, viz. Corroborants, as bark, olibanum, alum, elix. vitriol. bals. copaiv. &c. and if necessary, astringent injections and the cold bath.

GONORRHŒA Simple. See Gleet.
GONORRHŒA

GONORRHŒA Virulent. See Venereal

GOUT.

Symptoms. An acute pain in the joints, particularly of the feet. It is called regular when feated in the extremities, returns at stated periods, and gradually declines. Irregular, when the fits are uncertain and frequent, when the symptoms vary, and when the disease attacks the stomach, head, or other internal parts; chiefly affects the membranes, tendons, and ligaments; the pain like that of a dislocated bone, with a sensation as if warm water was poured on the part, succeeded by chilliness and slight fever: a breathing sweat comes on at the end of the fit, and the part swells; the fit returns at intervals, till the gouty matter is spent; the longer the intervals, the more severe the fit; but the symptoms are too well known, to need further enumeration in a work of this nature.

TREATMENT. Cordials are best during the sit; the body should not be costive; opiates may be occasionally given; the part should be wrapt in flannel, and relaxing or other topical applications, as the case may require; a temperate diet, exercise, friction, and the bath waters; some have ventured to blister the part with success.

GRAVEL. See Stone.

GUAIACUM Gum. See Gum.
GUAIACUM

GUAIACUM Wood.

Alterative, antirheumatic, antivenereal, in decoction, as Sarfaparilla.

GUM Ammoniac.

Pectoral, antispasmodic, gr. iv. ad pj. See Lac.

GUM Arabic.

Diuretic, emollient, in decoction ad libitum.

GUM Guaiacum.

Antirheumatic, alterative, antivenereal, Dj. ad 3j.

GUM Elemi.

Externally in digestives.

GUM Tragacanth.

As Gum arabic.

GUTTA SERENA.

SYMPTOMS. A blindness, though the eyes seem perfectly unaffected, the cause being in the

retina, or optic nerve.

TREATMENT. Blisters on the head, back, or behind the ears; bleeding if nothing forbids; valerian, castor, assafcetida, cinnabar, bark, volatile salts, millepedes, rosemary, or other nervous and attenuant remedies. Calomel, or chalybeates, if obstruction be the cause; emetics and cathartics discretionally; sternutatories should also be used.

HÆMORRHAGE external.

Apply balf. traumat. cobweb, dry lint, flour, agaric, alum, or the vitriolic styptics; if an ar-

tery

tery is injured, take it up, or apply the spunge tent.

HÆMORRHOIDS. See Piles.

HARTSHORN, Burnt.

Astringent, absorbent, Dj. ad zj. See Decoc-tion, white.

HARTSHORN Shavings. In jelly, corroborant, nutritive.

HEAD-ACH.

TREATMENT. If occasioned by plethora, use the pediluvium; bleed or cup, and purge: and if by viscid blood, give attenuants also; if by a foul stomach, an emetic; if by costiveness, purges; if it be nervous, castor, valerian, bark, spt. lav. c. spt. vol. arom. assafcetida; blister the back; and bathe the temples, forehead, &c. with æther, or spt. sal ammon. with a little roch alum in it; if a weak stomach, give bitters or other stomachics. Head-achs may arise from various other causes, which, when known, will indicate proper remedies.

HEART-BURN.

The common heart-burn, proceeding from acidity irritating the upper orifice of the stomach, magnesia, chalk, tabellæ, cardial. spt. vol. arom. or weak solutions of sal absent. remove it for the time; but to cure it, the stomachic corroborants must be given.

Heart-burn

Heart-burn may be occasioned by corroding humours of other kinds, and will not therefore be eased by alkalis; emetics, with plenty of chamomile tea or other liquid, to cleanse the stomach, are here proper; cathartics may also be useful, and these may be followed by stomachies; but water with gum arabic dissolved in it, will give occasional relief. Worms will also sometimes occasion this complaint; vermisuges are then proper.

HELLEBORE, Black.

Deobstruent, attenuant, alterative, gr. v. ad Aff. r more. See its Tineture and Extract.

HELLEBORE, White. Emetic, gr. v. ad off. sternutatory.

HEMICRANIA. See Head-ach.

HEMIPLEGIA. See Palfy.

HEMLOCK.

See its extract. In fomentations discutient and esolvent.

HEPATITIS. See Liver.

HERNIA, or Rupture.

TREATMENT. Return the intestines, if praccable, and confine it with a proper trus; offiveness must be avoided, as must likewise olent exercise, drunkenness, &c.

HERNIA Humoralis. See Testicles. HICCOUGH,

HICCOUGH, or Hiccup.

The common hiccup is usually removed by drinking a draught of any weak liquid, by holding the breath, and sometimes by swallowing dry bread.

For the hiccup or finguluts, which come on in the last stages of sever, &c. give julep. e mosch. or musk in substance, in large doses; volatile salt, castor or the like; to which, opium may be discretionally added. When it is a primary disease, sternutatories and emetics; and emp. stom. applied to the scrob.cordis have been useful.

HIERA PICRA.

Cathartic, deobstruent, Ass. ad Aj. or more. See Tinet. Sacra.

HONEY.

Pectoral, detergent.

HONEY, Egyptian, Detergent, destroys sungous slesh.

HONEY of Hellebore. Emetic, cathartic 3j. ad 3ff.

HONEY of Roses. Detergent, astringent.

HOOPING COUGH. See Cough.

HORSE RADISH.

Diuretic, deobstruent, ad libitum, or infused in water, wine, or spirit, zjss. ad ziij.

HYDROCELE,

HYDROCELE, HYDROCEPHALUS, See Dropfy.

HYDROPHOBIA.

It is of more consequence here to describe the symptoms in the dog than those of the patient. We may know the dog to be mad by his dull heavy look, endeavouring to hide himself; seldom or never barking; angry and snarls at strangers, but fawns on his owner; resuses all food, droops, hangs down his ears and tail, and often lies down

as if going to sleep; this is the first stage.

He now begins to breathe quick and heavy, shoots out his tongue, slavers, and froths at the mouth; looks half asleep; slies suddenly at by-standers; runs forward in a curve line; at length he knows not his owner, his eyes become thick and dim, and water runs from them; tongue of a lead colour; grows weak and faint; often falls down, then rises, and attempts to fly at something; grows mad and surious; the nearer to this state, the more dangerous the bite.

If the state of the dog cannot be obtained, he may be known to have been mad, or the disorder may be known to be this by the following effects, which will come on sooner or later: The bitten part begins to be painful, wandering pains come on gradually, with heaviness and uneasiness; disturbed sleep, frightful dreams, startings, spasins, sighing, anxiety, love of solitude; pains shoot from the bitten part up to the throat, with straight-

nefs,

ness, sensation of choking, and horror or dread at fight of water or other liquids, which is a suffi-

cient characteristic of the disease.

TREATMENT. Cut out the bitten part immediately, which is the only certain cure: cuppingglaffes should then be applied; cauterize and wash it daily with salt water, or ung. cærul. fort. keeping it open with escharotics; bleed, if the vessels be full; vomit with merc. emet. slav. give mosch. gr. xvi. cinnabar zsi. (to which camphire and pilfapon. ad gr. vii. may be added) at night, and a purge the next morning; bathe in the sea, or cold bath, and give a sudorisic; repeat these for a week daily, and three or four times at the next sull and change of the moon.

Ung. cærul. mit. applied externally, and turbith mineral given inwardly, so as to raise a salivation for several weeks, are said to be efficacious. The usual remedy known is now the Ormskirk medicine; but this has sailed in a variety of cases, and

has indeed been proved to be of no efficacy.

HYPOCHONDRIASIS.

SYMPTOMS are generally low-spiritedness (the disorder being chiefly in the imagination) heaviness, oppression, anddespondency; yet at times uncommon cheerfulness and flow of spirits; timidity, anxiety, fear, dread of dying, short cough, difficult breath, flatulency, pale urine, pains in the shead, odd fancies, spasms.

TREATMENT

tiously .

TREATMENT. Bark, and other tonics: neryous antispasmodics, as castor, valerian, assafoetida, &c. attenuants, as volatile falts and spirits; bitters and chalybeates if no fever; emetics, aperients, opiates, issues, and blisters discretionally; the cold bath and chalybeate waters, food light and easy of digestion, cheerful company, and gentle exercise.

HYSTERIA

In women, are fimilar to the hypochondriacs in men, but usually more violent, and attended with convulfive fits; which usually come on with oppression at the breast, difficult breathing, a semiof fomething rifing in the throat, which feems to threaten surrocation; convulsive motions; frothing at the mouth; laughing, and fometimes crying. May be known from a mere syncope by the pulse and breathing continuing, whereas in a fyncope they are not perceptible; it also comes on gradually, but a fyncope more fuddenly; in that, the fit is of short continuance, in this, usually much longer; with a colour in the face, which is not the case in a syncope.

TREATMENT of the diforder itself, as in the hypochondriaca. In the fit volatile fætids, finged feathers, &c. may be applied to the nostrils; and cold water and volatile fœtids given. The feet and legs may be placed in warm water; the pure cold air should be freely admitted; cold water may be sprinkled on the face and breast; and if a plethora be indicated bleed; otherwise cautiousis avoid it. In this disease, particular attention should be given to the state of the menses, &c.

This disease appears often under a variety of forms; for a more particular account than can here be given, Sydenham, and other authors may be consulted.

HYSTERITIS, or Inflammation of the Womb. See Womb.

JALAP.

Cathartic gr. xv. ad 3ss.

JAUNDICE.

SYMPTOMS. Yellowness of the whole skin, but chiefly the whites of the eyes; the urine also yellow; lassitude, inactivity, anxiety, sickness, oppression and difficult breathing, pain at the stomach, costiveness, hard, white, or grey stools, and bitter taste in the mouth.

TREATMENT. If the veffels be very full, bleed; then vomit, purge, and give medicines with foap; to which may occasionally be added, rhubarb, aloes, chalybeates or squills. Saline draughts if sever; opiates in case of pain; gentle emetics may be occasionally repeated, and the body should be kept open. Chalybeate waters, or water or cyder with a red hot iron quenched in it, may be used as common drink; gentle exercise, air, and cheerful company.

ILIAC PASSION.

SYMPTOMS. Acute pains in the bowels, oppression at the stomach, tension of the belly, vomiting of bile, and sometimes of sæces, great thirst and heat.

TREATMENT as in the inflammatory Colic, of which this is only a more violent kind.

INCRASSANTS, Bark and other Tonics, which fee.

INFLAMMATION Superficial. See Phlegmon.

INFLAMMATION of the bowels. See Colic and Iliac Passion.

INFLAMMATION of other Parts. See the respective Parts.

INFUSION Bitter.

Stomachic, corroborant ziss. bis die.

INFUSION Bitter, Purging.

Stomachic, corroborant, and opening; dose ziss. ad ziij. bis die.

INFUSION of Sena.

Purging zij. ad ziv.

INFUSION of Sena, with Lemons. Ditto.

IPECACUANHA.

Emetic off. ad zff. sudorific gr. sf. ad iij. with opium. Also in dysenteries.

2

IRON.

IRON. See Chalybs and its preparations.

ISINGLASS.

Corroborant, nutritive, in jellies.

ISSUES, to cut.

Pinch up the skin and fat with the thumb and forefinger, and divide them with a lancet so as to admit a pea; or you may apply a small blister or caustic on the part. The peas which it is afterwards dressed with may be smeared with basilicon or ung. ad vesic. if necessary; and dipped in blue vitriol water if sungus appears. The pea may be secured with adhesive plaister; if the issue instances, dress with ung. alb. &c. or bleed or purge.

ITCH.

SYMPTOMS. An eruption in the form of small pimples or pustules, chiefly about the joints, wrists, hams, the bend of the arms, the waist, and between the fingers; the pimples generally hard at first, afterwards watery, especially if rubbed or otherwise inflamed. An itching, which is greatest when warm, or in bed; it is also infectious.

TREATMENT. Anoint with fulph. viv. and exung. ung. cærul. or white precipitate and pomatum; hellebore, or fal ammoniac may occasionally be added; or wash with a weak solution of merc. corros. sub. a decoction of white hellebore root, or strong aq. veg. min. with salt. A quicksilver girdle may be worn,

but sulphur is the most safe and certain; fl. fulph. may be given inwardly; bleed and purge first in case of inflammation or costiveness.

JUICES, Scorbutic. zij. ad zvj. bis die.

JULEP of Camphire.
Diaphoretic, febrifuge, antispasmodic, diuretic, zj. ad ziij. every four, fix, eight, or twelve hours.

JULEP of Chalk.

Ab forbent, and in diarrhoeas, \$\frac{3}{2}\$ iff. ad \$\frac{3}{2}\$ iii. sepe:

JULEP of Musk.

Antispasmodic, (in singultus) 3j. ad 3 iij. frequenter.

JUNIPER Berries.

Carminative, diuretic, in decoction or infusion, ad libitum.

KERMES see Confection and Syrup.

KERMES Mineral.

Alterative gr. ff. ad gr. iij. bis die.

KIBES. See Chilblains.

KIDNEYS, inflammation of.

SYMPTOMS. Burning pain in the kidney, or kidneys; vomiting, eructations, the thigh feels numb, pain in the groin, ilium, and testicle of the same ide; frequent, but finall discharges of urine, which is red and high coloured, yet limpid and

watery in the height of the disease.

TREATMENT. Bleed; open the body with cathartics and clyfters; give emollient diuretics, with plenty of emulf. comm. or the like, for common drink; warm bathing; opiates may be

admitted in case of violent pain.

If the pain abates, leaving behind a fense of weight, with frequent shiverings, followed by heat, and whitish turbid urine, it denotes suppuration, balf. capiv. with nitre, and rhubarb, or other balfamic diuretics. The pain suddenly remitting, with cold sweats, intermitting pulse, hiccup, setid urine, and suppression of that discharge, are signs of mortification.

KING's EVIL. See Scrophula.

LABDANUM. Used in warm plaisters.

LABOUR. See Child-birth.

LAC AMMONIACUM. Asthmatic, antispasmodic zj. ad ziij.

LAC, Tineture of. Dentrific.

LAVENDER, Infusion of. Cephalic, nervous.

LEAD.

Externally deficcative, repellent.

LEMON Juice. Cooling, antiseptic zj. ad zss.

LEMON

LEMON Peel.

Stomachic, in infusion 3j. ad 3iij.

LETHARGY.

This may be confidered only as a leffer degree of Apoplexy, and should be treated accordingly. The chief symptoms are sleep, or great drowsines.

LEY, or LIXIVIUM of Soap.
Antacid, diuretic, lithontriptic, gut.x. ad zj.bis die.

LEY, or LIXIVIUM of Tartar. Antacid, diuretic, gut. xv. ad 3j.

LIENTERY.

In this disease the aliment is discharged with out little alteration, and the body wastes. The Treatment as in the Cæliac Passion.

LIME. See Water.

LINIMENT, Arcaus's. Digestive, incarnative.

LINIMENT Mercurial. See Ointment
Mercurial.

LINIMENT Saponaceous. Discutient, strengthening.

LINIMENT Volatile. Discutient, resolvent, repellent.

LINIMENT White.

Healing, softening.

LINIMENTUM

LINIMENTUM Tripharmacum.

Drying, healing.
LINSEED Infusion of.

Pectoral, diuretic.

LITHON TRIPTICS.

Lix. sapon. gut. x. ad zj. Soap zf. ad zij. Sal ammon. vol. gr. v. ad xv. Spt. sal marin. gut. x. ad xx. or more. Lime-water (oysterthell) ad libit.

LIVER inflamed.

SYMPTOMS. Great pain in the part, shooting; up towards the throat and shoulder; vomiting, dry cough, lassitude, fever, tension of the hypochondrium; eyes, ikin, and urine, sometimes; yellow; costiveness, difficulty of breathing.

TREATMENT. Bleed as largely as the pulse: will bear; blifter the part, purge, give cooling; febrifuges and diuretics; direct cooling emollient: clysters, fomentations, and poultices, and pediluvia...

If a tumour appears, forward and open it, if

nothing forbids.

In the East-Indies they cure this disease by: falivation, bleeding and purging being premised.

If a Schirrus succeeds, soap, rhubarb, mercurials, and chalybeates will be proper.

LOCHIA. See Child-birth.

LOCKED JAW.

SYMPTOMS. A rigid contraction of the muscles which raise the lower jaw, whence the jaws. continue continue violently closed, with great pain.

TREATMENT. The best remedy is large and repeated doses of opium. Blisters, purges, clysers, and antispasmodics may also be prescribed.

LOGWOOD, Infusion of. Astringent zij. ad ziv. See Extract.

LOTION, Soap. For cutaneous deformities.

LUES VENEREA. See Venereal Disease.

LUMBAGO.

A kind of rheumatism in the loins or small of he back, without any nephritic symptoms. Anirheumatic remedies prove most efficacious. See *Rheumatism*. Strengthening or stimulating plaisters, or even blisters to the part, are also seful.

LUXATIONS.

TREATMENT. Extend the limb or part, fo hat the head of the bone may be free from obstales; then reduce the bone into its natural situation, and apply proper bandages, not too tight, to eep it so till the ligaments have recovered their one. If there be inflammation, bleed, give laxtives and cooling remedies, foment and poultice. Lest and proper diet should be advised.

MADNESS.

The fymptoms are too well known to need numeration. Emetics, cathartics, blifters, occaonal bleeding, antispasinodics, opiates, and low diet.

diet. These may be varied according to the nature of the complaint (that is, whether it be Melancholy or the contrary) and to the symptoms that arise in either. Dr. Simmons, who has made trial of a great variety of remedies at St. Luke's Hospital, has often seen very remarkable good effects, and even cures performed, by large doses of camphire in maniacal cases.

MAGNESIA.

Antacid, cardialgia, opening Dj. ad 3ij.

MEASLES.

SYMPTOMS. Chilliness, shivering, pain in the head, fever, fickness, and sometimes vomiting; cough, heaviness, swelling and inflammation of the eyes, and discharge of watery humour from them, and also from the nostrils. The third or fourth day, an eruption like flea-bites appear in the face, neck, and breast, and soon after in the body and limbs; they rife above the skin, but do not suppurate. The fever and other symptoms do not, as in the Small Pox, &c. abate on the appearance of the eruption, which continues about three days, then dries away, the fkin peeling of; but the other symptoms remain and even increase, especially the cough, which is also attended generally with difficulty of breathing; and oppression at the breast.

TREATMENT. Cooling and aperient febrifuges; gentle diaphoretics, if necessary; bleeding, if

nothing forbids; pectorals for the cough; opiates occasionally at night; blifter, if the cough be obstinate; and bathe the feet in warm water. Gentle physic should be given as soon as the

eruption disappears.

It has been proposed to inoculate the Measles; but Dr. Simmons, in his differtation de Rubeola, observes, that the different attempts in this way, hitherto have proved ineffectual; Dr Home's experiments, as related in his Medical Facts, not having been found to answer when repeated by others.

MECONIUM retention of, in Infants; (called also Colica Meconialis.)

TREATMENT. Give magnefia or fyr. rof. with ol. ricini, or ol. amygd. ol. anifi, or other pil in case of wind, and if necessary, emollient lysters. If purging, give creta, or other absorpents, &c. If vomiting, fyr. e mecon. given with aution in small doses.

MENSES, immoderate flux of.
TREATMENT. Tinct. rolar. tinct. styptic, lum, bark, pulv. e fuccin. c. opiates. In deferate cases, small doses of vitriol. cærul. or tinct. iturn. but with great caution. Bleeding, aperints, aftringent fomentations to the part, as vinear, alum, or the like; rest, and proper mild diet.

MENSES obstructed.

TREATMENT. In phlegmatic habits, purge ith tinct. facra pil. rufi. or other aloetics, and

give chalybeates. In plethoric habits, bleed, avoid chalybeates, and give pulv. e myrrh. c. castor, or the like, with tinct. melampodii, aloes, oil or extract of savine, pennyroyal, and the like. Bathing the feet, somenting the belly, and sitting over the steams of hot water are good in both habits; attenuants in case of viscid blood; emetics are often efficacious, as are sometimes small doses of mercurials; exercise, and if the appetite be vitiated, proper stomachics. Of late, electricity has been sound efficacious in cases of this sort, by drawing sparks, and sending gentle shocks through the pelvis.

MERCURIAL Pills. See Pill.

MERCURIAL Plaister, Ointment, &c. See Plaister and Ointment.

MERCURIUS dulcis. (Calomel.)

Alterative, antivenereal gr. j. ad v. omninocte. (Also vermisuge, deobstruent, and for salivation) Cathartic gr. v. ad xv.

MERCURY Alkalized.

Alterative, vermifuge gr. v. ad zff. bis die.

MERCURY Calcined.

Alterative, antivenereal gr. ff. ad ij. omni noste.

MERCURY Corrosive sublimate.

Alterative, antivenereal gr. 1/6 ad gr. j. bis die.

MERCURY Red corrosive. Escarotic. MERCURY

MERCURY precipitate, White. Antipforic, &c. in liniments.

MERCURY Emetic, Yellow. Gr. ij. ad iv.

MEZERION.

Antivenereal, alterative, in decoction, dose zij. ad iv.

MILIARY MILK See Fever. See Fever.

MILLIPEDES.

Diuretic, attenuant, deobstruent, number ij. ad v. or more alive; or in powder gr. iij. ad xv.

MISCARRIAGE, See Abortion.

MITHRIDATE.

Alexipharmic, anodyne, Dj. ad 3iff.

MORTIFICATION. See Sphacelus.

MUSK.

Antispasmodic gr. iij. or more. See Julep.

MYRRH, Why was

Deobstruent, antispasmodic gr. iij. ad xij.

NEPHRITICS.

Gum arabic ad libit. Spt. nitr. dulc. gut. xx. ad zj. Sal diureticus əff. ad 3 j. Opiates. See Diuretics.

NEPHRITIS. See Kidneys.
NERVOUS

NERVOUS Complaints. See Hypochondriac

and Hysteric Difeases.

The class of medicines called Nervous, with corroborants, air, and proper regimen, are most useful in these cases.

NERVOUS Fever. See Fever.

NERVOUS Medicines.

Pil. gum gr. v. ad bj. Aslafætida, and the other gums in like doses. See also Bark, the Chalybeates, Cephalics, Tonics, and Antispasmodics.

NIPPLES fore.
TREATMENT. Apply oil of wax, cream, mucilage of gum Arabic, ung. alb. fuet, or the like. If they are moift, ung. alb. with fome drying powder, as starch, bol. armen. &c.

NITRE.

Cooling, febrifuge, diuretic gr. v. ad Dj. OIL of Almonds. Opening, pectoral; out-Olives. wardly, softening, relaxing.

OILS Essential.

The virtues of the plants they are drawn from, in doses of gut. j. ad iij. on sugar, or otherwise.

OILS by Decoction and Infusion. Softening and relaxing externally applied.

OIL of Palm.

Outwardly for cramps, weaknesses of nerves, and chilblains.

OIL

. OIL of Wax:

Externally to chaps, &c.

OINTMENT Bafilicon.

Digestive.

OINTMENT of Elder.

- Marshmallows.

Cooling, emollient.

OINTMENT of Gum Elemi. Digestive, incarnative.

OINTMENT Mercurial.

For falivation; repellent, discutient, for the itch, vermin, &c.

OINTMENT of Precipitate.

- Sulphur.

For the itch, &c.

OINTMENT Saturnine.

Drying, repellent.

OINTMENT of Tar.

For feald heads, digestive, &c.

OINTMENT of Tutty.

For the eyes.

OINTMENT White.

Healing, foftening, cooling.

OLIBANUM.

Corroborant, gr. v. ad Dj. OPIUM.

Anodyne, antispasmodic, &c. gr. 4 ad ij.

3 2 OFOPANAX

OPOPANAX.

Nervous, antispasmodic, gr. v. ad xv.

ORANGE Juice, As lemon, but weaker.

OXYMEL Simple. Antiseptic, cooling, detergent.

OXYMEL of Squills.

Expectorant, diuretic 3 st. ad 3 j. Emetic 3 ij. ad 3 st.

OYSTER-SHELLS.

Absorbent gr. xx. ad 3j.

OYSTER-SHELLS calcined. (Their water) Lithontriptic ziv. or more.

PALPITATION of the Heart.

TREATMENT. In proper constitutions bleed repeatedly; but where the disorder proceeds from relaxation, the bark and corroborants. The nervous tribe, musk, volatile salts, &c, are good; as are also blisters.

PALSY, or PARALYSIS.

Symptoms. A loss or diminution of the motion or feeling, or both, or some part or parts of the body. If it happens to all the parts below the head, it is called *Paraplegia*; if to one side of the body, *Hemiplegia*; and if to a particular part *Paralysis*.

TREATMENT. Emetics, blifters kept open, and strong clysters; nervous, attenuating, and stimulating

stimulating medicines, as valerian, castor, mustard, sal c. c. horse-radish, &c. And mustard, or other warm stimulating application rubbed into the parts; and to that part of the spina from whence the nerves serving the part issue. Electricity and dry frictions are good, as are also chalybeate, and Bath or Bristol waters.

PEARLS.

Absorbent Dj ad zj.

PECTORALS.

Elix. pareg. zss. ad zj. Pulv. e tragac. c. Dj. ad zj. Pil. de styrace gr. v. ad Dj. Oxymel scillitic. zss. Troch. bechic. decoct. pect. Ad libitum.

PEEL. Sec Bark.

PERIPNEUMONY, or inflammation of the Lungs.

SYMPTOMS. Difficult breathing, with oppression and seeming load at the breast; the breath hot, cough, sever, redness in the sace, pulse sometimes hardly perceptible, but after bleeding, trong, but unequal. Differs from a *Pleurisy* in the cough being more moist, the pain less acute, for the pulse so strong.

TREATMENT. Bleed repeatedly, if necessary; nd give nitre, with spermaceti, or ol. amygd. sal. c. gentle antimonials, or the like. Cooling mollient clysters are useful, as are also blistering

nd cupping.

PERIPNEUMONY,

PERIPNEUMONY, Bostard.

SYMPTOMS. Heat and cold alternately; pain and giddiness of the head, when the cough is most troublesome; vomiting, quick, difficult breathing, and wheezing; pain of the breast; urine turbid, with red sediment. Differs from a true Peripneumony, in having neither extraordinary heat, pain, thirst, or sever; and by its usually seizing those of a relaxed and gross habit; prevailing most in moist foggy weather, contrary to the other.

TREATMENT. Bleed not, unless the pulse be very full, and comatose symptoms appear; blisters, emetics, laxatives, emollient clysters, attenuants and diluents; pectorals for the cough, and opiates

if necessary.

PHILONIUM London. Carminative, anodyne off. ad 3 ff.

PHLEGMON, or superficial Inflammation.

SYMPTOMS. Heat, pain, tension, redness, and throbbing in the part; fever; that part proceeding to suppuration or solution, gangrene, or scirrhus.

TREATMENT. Remove any perceivable external cause; bleed, purge, and give diaphoretic febrifuges; apply emollient fomentations and poultices to the part. If matter forms, cease evacuation, and forward the suppuration; when ripe, open the abscess and digest it off. If gangrene appears, treat as directed under that article.

PHRENSY.

PHRENSY, Phrenitis, or Inflammation of the Brain.

SYMPTOMS. Constant delirium and sever, ifficult breathing, violent pain in the head; reducts of eyes and face, pulse quick, hard, and small; ongue black and dry, urine thin and transparent, udden startings, picking the bed-clothes. Inown from a common sever, &c. by the delirum preceding the sever.

TREATMENT. Bleed largely; stimulating lysters, blisters, the pediluvium, and sinapisms to he feet; spirituous embrocations to the head, rst shaved. Nitre, sal c. c. or other attenuants, urges, and diluents; promoting the hæmorrhoi-

al flux is of great ufe.

PHTHISIS. See Consumption.

PHYMOSIS and PARAPHYMOSIS.

SYMPTOMS. The first is a closing of the preuce over the glans penis, so as not to be drawn ack; the last, a constriction of it behind the lans, so as not to be drawn forward over it.

TREATMENT. Foment, poultice, bleed, and urge; vomit if necessary; and give cooling feb-fuges. If these fail, proper incisions must be

ade.

ILES, or painful Tumours in the Anus or Rectum.

TREATMENT. Bleed, if necessary; and give lphur, or æthiops mineral, nitre, and elect. nitiv. opiates if they are very painful; and if they

they are external, foment them with warm emold lient liquors, or anoint them with ung. famb. on the like; if they bleed much, apply gentle aftringents; if much swelled and inflamed, scarify, or apply leeches to them. Ball. sulph. is very proper to apply to the Piles, and the ung. cærul. has also been found useful. They may more expeditiously be driven away by liniments made of powder or galls, or sugar of lead, or by washing them with solutions of white or blue vitriol. But as the hæmorrhoidal discharge is generally salutary, caustion must be used in repelling them.

PILL Aromatic.

Aperient zss. and zj. antirheumatic, deobstruent gr. v. ad xii. bis die.

PILL Colocynth.

(Both) Cathartic Dj. ad Dij.

PILL Ecphractic. Emmenagogue gr. v. ad zsf. bis die.

PILL Gummous.

Hysteric, nervous, &c. gr. v. ad 3ss. bis die.

PILL Mercurial.

Alterative, antivenereal gr. v. ad 9j. omni nocte

PILL Rufus's.

Deobstruent gr. v. ad Dj bis die. Opening Dj. ad 3ss.

PILL Soap

Diuretic, icteric, anodyne gr. v. ad Dj.

PILI

PILL Storax.

Pectoral, anodyne gr. iv. ad 9j. at night.

PILL Squill.

Asthmatic, pectoral, diuretic gr. v. ad xij.

PIMPLES.

TREATMENT. Wash them with a solution of sch. saturn. or vitriol. alb. If these sail, try spt. in. camph. a solution of sal tartari, or white preipitate mixed with pomatum. If they proceed rom soul blood or a scorbutic cause (as it is termed) reat them with alteratives, as directed for the Land Scurvy; with now and then a cooling urge.

PLAGUE.

SYMPTOMS. Shivering; pain in the head, ack, and ftomach; fickness and vomiting; defondency, anxiety, difficult breathing, wildness of ountenance, high fever, faintings, hiccups, and atching of the tendons. Tumours at length ppear in the groin, arm-pits, or behind the ars; and then the fever abates, but returns if the umours fink. Fætid breath and sweat; livid pots, fornetimes broad, and fuddenly difappearing. Carbuncles arise in the worst state of the disease, which feldom suppurate kindly, but appear as red iery circles; are fmall tumours with corrupted lesh underneath, and are attended with great tching. The fymptoms, however, vary accordng to the constitution and state of the air, but hese are the most general.

TREATMENT:

TREATMENT. The tumours are critical, and should be encouraged by ripening cataplasms and proper cordials; the carbuncles should be poulticed with cataplasm e cymino, or the like; and iff they appear gangrenous, scarify, and dress them with ol. terebinth, and hot digestives, as in Mortifications.

The disease itself should be treated with diaphoretics, antiseptics, cordials, antiphlogistics, or other remedies, according to the symptoms; but in general, camphire, nitre, snakeroot, and bark, &c. and emetic being first given, seems most eligible in

this dreadful disorder.

N. B. The vapour generated by nitre, especially if the deflagration be made with odoriferous

ingredients, is faid to prevent infection.

Inoculation has lately been recommended in the *Plague* by Dr. Samoilovitz, a Ruffian phyfician; but if the disease (as seems to be the case) is liable to attack more than once, this can be of no use.

PLAISTER Common.

Healing, drying.

PLAISTER Common, with Gums, Suppurative.

PLAISTER Mercurial.

Discutient, resolvent.

PLAISTER Mucilage.

Softening, and for corns.

PLAISTER

PLAISTER Red Lead.

Drying, healing.
PLAISTER Soap.

Resolvent:

PLEURISY.

SYMPTOMS. Chilliness, shivering, succeeded by heat, thirst, and restlessines; a violent acute pain on one fide near the ribs, extending towards the shoulder-blades, back, the breast; worst when the affected side is lain on. Difficult breathing, redness of the cheeks, nausea, and dry cough; the matter spit up if any, yellowish or bloody.

TREATMENT as in the Peripneumony; but repeated bleeding, laxatives, and cooling attenuants, as fal c. c. with nitre are chiefly to be

relied on.

POISONS.

TREATMENT. When these are taken, if the time has been but short, give a strong emetic of white or blue vitriol, drinking plentifully of warm water to wash the stomach; oily clysters should also be given repeatedly, as should likewise cathartics. If corrosive sublimate was the poison, give folution of any fixed alcaline falt. If opium, or other narcotic, give acids (as vinegar) with water or broth; apply blifters; and if necessary, bleed. Care must be taken to continue these operations a fufficient time; afterwards the fromach and bowels should be healed with proper balsamic remedies, broths, &c.

POMEGRANATE

POMEGRANATE SHELL. See Bark. The juice cooling, antiseptic.

POPPIES. See Syrup of Poppies.

POWDER of Amber, compound.
Astringent, corroborant gr. viij. ad 3ss. bis die.

POWDER Antilyssus. For bite of mad dog 3ss. omni mane.

POWDER of Arum root, compound. Stomachic, antirheumatic Dj. ad zj. bis die.

POWDER Bezoardic.
Absorbent, alexipharmic Dj. ad zj. at night.

POWDER of Bole, compound, Astringent Hst. 3st. bis die.

POWDER of Ceruse, compound.

Cooling and astringent, in collyriums for the eyes.

POWDER of Contrayerva, compound, Sudorific Dj. ad zj.

POWDER of Crabs Claws, compound.

Absorbent, febrisuge, Dj. ad zj. frequenter.

POWDER to promote Delivery. Dj. ad Zj.

POWDER of Scammony, compound. Cathartic Hi. ad Hij.

POWDER

POWDER of Sena, compound. Cathartic 3 sf. ad 3 j. or more.

POWDER of Tin. Vermifuge Dj. ad Z j. bis die.

POWDER of Tragacanth, compound. Pectoral 3j. ad 3j. frequenter.

POX. See Venereal Disease.

POX, Chicken; or SWINE Pox.

SYMPTOMS. Chilliness, heat, and the usual precursory symptoms of sever; but milder than in the small pox. The eruption appears about the third day. They rise like the small pox, but may be known from them by the matter being more like water than pus, by little vesicles, or bladders filled with water on the top of such pushules as are whole; and by their turning about the fifth day; which the small pox does not till at least the seventh; and they leave no marks behind them.

TREATMENT. Cooling febrifuges, as nitre, with magnefia, antimonials, pulv. contray. and faline draughts, &c. and diluents; and afterwards two or three purges.

The Swine Pox is only a larger fort of

Chicken Pox.

POX, Small. See Small Pox.

PROCIDENTIA, vel PROLAPSUS ANI, or bearing down of the Rectum.

TREATMENT.

TREATMENT. Corroborants, as bark, tinct. rofar. &c. Laxatives, clyfters, and aperients, if costive; clysters of tinct. rosar. or other bracers, and a proper bandage; chalybeates and cold bathing. N. B. It there be inflammation, emollient fomentations and poultices. part should be gently reduced, and kept up with a bandage; rest and an horizontal posture are proper.

Prolapsus Uteri. TREATMENT as in the preceding case. Pessaries are also proper to be worn.

PUERPERAL PUTRID | Fever. See Fever.

QUICKSILVER.
Vermifuge, opening zff. ad zij. or more, in the morning.

QUINCY.

SYMPTOMS. Inflammation of the throat; with pain, great heat, tumour, redness, difficulty in swallowing, and fever. The uvula sometimes relaxed and inflamed; the face red.

TREATMENT. Bleed largely, or according to exigency; open the body; blifters to the back, and near the part; gargle with fpt. fal. ammon. and decost. pect. strong of the former; poultice the part with bread and milk, to which camphire may be added, and the volatile liniment may also be applied. If it suppurates, cease to evacuate; and inhale emollient steams. When ripe open the abscess; afterwards gargle with tinct. rosar. and tinct. myrrh.

RATTLESNAKE ROOT.

Diaphoretic, antiseptic gr. v. ad Dj. Also in antiseptic gargles.

RESIN of Aloes.

Cathartic Dj. or more. Deobstruent gr. v. ad Aff.

RESIN of Bark.
Corroborant, febrifuge, &c. gr. v. ad Dj.

RESIN of Guaiacum

Antirheumatic, alterative, deobstruent, gr. v. ad xv.

RESIN of Jalap.

RESIN of Scammony. Cathartic gr. v. ad Aj.

RHEUMATISM Acute.

SYMPTOMS. Rigor, shivering, fever, thirst, costiveness, wandering gnawing pains, with fwelling and redness in the part affected. The

pain worse towards night, and in bed.

TREATMENT. Bleeding; medicines with fall c. c. nitre, camphire, guaiacum, antimonials, or the like; laxatives, and opiates occasionally. The part may be rubbed with lin. fapon. camphire, and oil; lin. vol. &c. Warm fomentations are good; the part may be covered with

flannel, diluents should be freely given, and a diaphoresis kept up. When the urine deposits a copious sediment, the bark may be given; it may be joined with some of the above, and opium may be occasionally added.

RHEUMATISM chronic.

Differs from the acute, in that it is not attended with fever or inflammation in the part; and chiefly attacks those advanced in life, contrary to the other.

TREATMENT. If necessary, bleed, and purge with elect. e scammon. sena, &c. Pulv. doveri, or other anodyne sudorisic, may be given. Calomel in small doses, with vin. antim. has been sound efficacious; as hath likewise kermes mineral, with camphire, about three grains of each. Bark and sarsaparilla are given by some; but gum gualiacum with volatile salts, the guaiacum with an opiate, or sal c. c. with nitre, and julep e camph. according to circumstances, are generally successful. Warm embrocations to the part, and slannel are also serviceable; cold bathing, where not improper on other accounts; a slannel shirt worn next the skin, is a good preventive of a return. N. B. Mustard has been effectual, taken inwardly, and rubbed into the part. Electricity, friction and exercise of the part, are useful.

RHUBARB.

Cathartic Dj. ad zj. Stomachic gr. iv. ad zff. RHUBARB

RHUBARB toassted.
Astringent Hs. ad Hj.

RICKETS.

SYMPTOMS. The head large, the fontanellar keeps long open, the face full and florid, the joints knotty, and the bones protuberant, causing incurvation and distortion; the ribs protuberate, and become crooked; the belly swells; cough, and pulmonary disorders succeed. The child (for children only are the subjects of this complaint) moves weakly, and waddles; usually appears about the eighth month, and continues till the fixth year; ariseth from bad nursing; the under-standing is very early.

TREATMENT. Cold bathing; friction; bathing the back with spirituous embrocations, or warm balsamic liniments; strengthening plaisters; gentle vomits of ipecacuanha; and purges with rhubarb and æthiops mineral; bark, chalybeates,

air, and exercise.

RING WORMS. See Tetters.

ROB of Elder. Pectoral, opening zj. ad ziij.

ROSES, red.

Astringent gr. v. ad 3ss. See Tincture.

ROSEMARY.

Cephalic, carminative, &c. in infusion.

RUE,

RUE.

In infusion against sits and spasmodic complaints; also alexipharmic and antipestilential; juice in clysters, vermisuge.

RUPTURES. See Hernia.

SAFFRON.

Cordial gr. iij. ad pj or in infusion.

SAGAPENUM.

Nervous, hysterie, &c. gr. v. ad Dj. bis die.

SALT, alkaline fixed.

(As fal absinth. fal tartar, &c.) antacid, attenuant, diuretic gr. iij. ad xv.

SALT, alkaline volatile.

(As fal c. c. fal. ammon. vol. &c.) antacid, attenuant, diaphoretic gr. iij. ad xv.

SALT of Amber.

Aperient, diuretic, hysteric gr. iv. ad 9j.

SALT Ammoniac.

Diuretic, febrifuge, attenuant, diaphoretic gr., v. ad Dj. or more.

SALT Cathartics

(As Glauber's, Epsom, Rochelle, &c. 3ss. ad 3ij.)

SALT Diurerie.

Attenuant 3j. ad 3ij. Aperient 3ij. ad 3ff. SALT

SALT of Steel.

Attenuant, decbstruent gr. ij. ad off.

SARSAPARILLA.

Alterative, antivenereal, in decoction (zij. ad thij.) ziv. ad this die.

SASSAFRAS.

Alterative, antiscorbutic, &c. dose as in Sarsa-parilla.

SAVIN.

Emmenagogue, in infusion zij. ad ziv.

SCALDS, or Burns.

TREATMENT. Apply ol. lini f. igne alone, or with spermaceti or wax dissolved in it; aq. yeg. min. is also good; afterwards cerat. epulot. or other cicatrizer. If inflammation renders it necessary, bleed and purge.

SCALD-HEAD.

TREATMENT. Rub the part with ointments made of fulphur, pitch, ung. nicotian, or the like; the ung. cærul. has been used with success; bathing with salt water, and give mercurials, antimonials, or other alteratives; blisters kept open, and issues, are proper.

SCAMMONY.

Cathartic off. ad Dj.

SCARLET Fever. See Fever.

SCHIRRUS, or hardened Glands.

TREATMENT.

TREATMENT. Endeavour to disperse it by mercurial unction, ext. cicutæ, or mercurials, with sarsaparilla, &c. and occasional purges. If it inclines to suppurate, treat it as directed in abscesses, &c. If it neither disperses nor suppurates, extirpation is the only cure, where it can be done; otherwise treat it with coolers, and occasional anodynes.

SCIATICA.

SYMPTOMS. A fixed pain in the hip, like the rheumatism. Called also the Hip Gout and Ischias.

TREATMENT as in Lumbago.

SCORDIUM.
Astringent. See its Preparations.

SCROPHULA.

SYMPTOMS. Schirrous unequal tumours, which do not readily suppurate; affecting the glandular parts, but chiefly on the sides of the neck.

TREATMENT. Purge occasionally with jalap and calomel; and give alteratives, as calomel, with sulph. aur. ant. æth. antimon. or some other; millepedes are good; spong. ust. with rhubarb; sea-water internally before suppuration, and bark and sea-bathing after they soften; exercise is good; gross and high seasoned diet must be avoided; vegetable acids, especially Seville oranges, are proper.

SCURVY.

SCURVY.

SYMPTOMS. Heaviness, lassitude, low spirits, offensive breath, tender gums, sallow bloated countenance, hæmorrhages from the nose and nouth, difficult breathing, swelling of the legs, rellow, purple, or livid spots on the skin; tumours in the limbs; contraction of the tendons of the sam; other symptoms occur, but they differ in lifferent subjects, as do likewise those above-menioned; and the disease is in general sufficiently thown. It is, however, distinguished into Sea and Land Scurvy.

TREATMENT. In the Sea Scurvy, the juices re exposed to putridity from the use of animal bod, and moist air; antiseptics therefore will be roper, as vegetables, vegetable acids, fruit, cyder, cc. Insusion of malt, in defect of these, or quors made of molasses or sugar. Bark, and lix. vitr. acid. good. Liquids impregnated with xed air for common drink, and wholesome air. Thalybeates have been found serviceable, especially

then joined with the bitters, or bark.

The Land Scurvy (improperly fo called) is ther a cutaneous disease; scurfy or scabby erupons appear, either partially, or more universally;

ften with itching and heat.

The antimonial alteratives, with gentle mercuals, are often efficacious; aq. calcis c. succeptudic, or the like, may be used with them ream of tartar and flowers of sulphur are good he parts may be anointed with saturnine lini-

ments, with a little white precipitate if necessary.

SCURVY GRASS.

Antiscorbutic, the juice or insusion zij. ad ziv. bis die.

SENEKA. See Rattlefnake root.

SENNA.

Cathartic 3 j. ad 3 iij. or in infusion.

SETON, to make.

Pinch up the skin and fat with the finger and thumb, and run a proper needle through it, armed with a skain of silk or thread smeared with digestive ointment; which must be shifted a little every day, so as to keep both orifices running.

SHINGLES. See Tetters.

SINGULTUS. See Hiccups.

SKIN, deseases of. See Cutaneous Diseases.

SLATE, Irish.

An internal vulnerary and aftringent, 3 j. ad 3 ij. in spruce beer.

SMALL-POX.

SYMPTOMS. Chillines, shivering, heat, pain in the head and back, sickness and pain at the stomach, and sometimes vomiting; sever, costiveness, convulsive fits sometimes in children, which is a good symptom. On the third or sourch day the eruption appears, like slea-bites, in the face, breast, arms, body, and feet successively;

pains .

pains and foreness in the throat. The eruption being compleated, the fever goes off or abates. The eruptions fill, or maturate like boils, and the skin between them is also red; the eye-lids, face, hands, and feet, swell successively; the pocks from red, grow whitish and smooth in the body, and yellow and rough in the face, full of a white matter. About the seventh day from the eruption, (sometimes the ninth) they dry, or turn as it is called, and scale off, leaving red

marks, and fometimes pits behind.

TREATMENT. If the pulse be full and strong at the beginning, bleed; and if necessary, or the period be sufficiently early, purge, and give an emetic. During the eruptive sever, cooling sebrifuges, with laxatives, or clysters occasionally; but if the symptoms require it (as low pulse, &c.) gentle cordials and diaphoretics. After the eruption, little is generally required but a gentle anodyne at night, proper diluents, and if necessary, opening clysters, &c. If purging comes on, check it with caution; if delirium or sever, apply blisters; and if the pock sinks, give bark, with proper diaphoretics and cordials; especially if malignant or putrid symptoms, or petechiæ appear. Spt. vitrioli dulcis is also good in these cases, as are sinapisms to the sect.

In the confluent kind, all the symptoms are worse than in the distinct. The eruption makes its appearance a day or two later, and does not

come

come to its height till the eleventh or fourteenth day after; the fever also continues after the eruption appears, and a salivation comes on, which must by no means be checked. The patient requires to be supported more in this case than in the other; care, however, must be taken in both kinds to avoid too much heat, or a heating regimen; and pure, and even cool air, where no objection appears, is salutary. If the pustules are opened, and the matter let out, so much the better; especially if danger of their sinking or striking in appears. Purging is esteemed proper after this disease.

To inoculate for the Small-Pox, take a little matter from a puffule, after the eruptive fever, on the point of a lancet, and insert it in the arm (by making a small puncture or two) between the true and scarf skin, which may afterwards be pressed down with the flat side of the lancet. Three or sour days afterwards the punctures appear inslamed; in about three days more the precursory symptoms come on, and at length the eruption.

Previous to the inoculation, a mercurial purge or two may be administered, if the patient is of a full or gross habit; and he should also be restrained to a milk and farinaceous or vegetable diet, avoiding especially high seasoned food. It is found best to keep him out of bed, and even now and then in the cold air. The symptoms are

generally

generally so favourable, that little affistance is required from medicine, except a few mercurial purges after the disorder goes off.

SOAP.

Diuretic, icteric, lithontriptic, attenuant Dj. ad Z ij. bis die.

SOAP Leys. See Lixivium.

SORE THROAT, putrid. See Angina maligna.

SPASMS. See Convulsions.

SPECIES Aromatic.

Gr. v. ad xv.

SPECIES of Scordium.
Astringent gr. v. ad pj.

SPERMACETI.

Pectoral, balfamic, vulnerary Dj. ad 3 j. Externally healing, softening.

SPHACELUS, or complete Mortification.

SYMPTOMS. This succeeds a Gangrene (which see) if it cannot be stopt; the part looses all sensation, and yields an intolerable smell.

TREATMENT as in Gangrene. When the separation is advanced, amputate the limb, or cut out the part, if it can be done with safety; persisting in the use of the bark, &c. till a complete digestion is procured.

SPIRIT

SPIRIT of Hartsborn.
Attenuant, cordial, sudorific gut. x. ad lx.

SPIRIT Mindererus's.

Attenuant, cooling, diaphoretic 3 ij. ad 311.

SPIRIT of Nitre, Sweet.

Diuretic, febrifuge, carminative gr. x. ad 3 st.

SPIRIT of Turpentine.

Diuretic, &c. gut x. ad xl.

SPIRIT of Vitriol, acid.

Cooling, aftringent, antiseptic, antalkaline, gut. v. ad xx.

SPIRIT of Vitriol Sweet.
Febrifuge, diuretic, &c. gut. v. ad xx.

SPIRIT Volatile aromatic. As spirit of Hartshorn; also cephalic.

SPIRIT Volatile fætid. Antispasmodic, &c. gut. ad xxx.

SPLEEN. See Hypochondriacal and Hysteric Diseases.

SPRAINS. See Strains.

SQUILLS.

Pectoral, diuretic gr. iij. ad viii. Emetic off.

STEEL, Ruft of.

Deobstruent, attenuant, tonic gr. v. ad off. See its Preparations.

STINGS,

STINGS, Venomous. TREATMENT as in Bites.

STOMACH, Inflammation of.

SYMPTOMS. Heat, pulsation, and acute pain in the part, which is increased by swallowing; tension at the pit of the stomach; anxiety; retching; hiccup; small, weak, intermitting pulse.

TREATMENT. Bleeding, emollient liquids, and clysters; fomentations; gentle opiates; blisters; mucilage of gum arabic; with small

doses of nitre occasionally.

STOMACH, pain in. See Heart-burn.

STONE AND GRAVEL.

SYMPTOMS. Pain in the back and loins, or the region of the bladder; nausea; vomiting; strangury; sometimes bloody urine; costiveness; retraction of the testicle of the affected side; numbness down the thigh and leg; the pain in the kidneys abate when the calculus is got into the bladder, and the urine becomes turbid.

TREATMENT. Bleed, if no objection appears; draughts with ol. amygd. and large quantities of gum arabic, with manna and opium added occafionally. Pil. sapon. good as an anodyne at night. As solvents, the lixiv. sapon. from x. to xxx. or lx. drops in veal broth twice in a day. Soap, oyster-shells, and lime-water are also good; and in some cases, the sal vol. c. c. or spt. sal. marin. Water impregnated with fixed air, and the effer-

1 2

vescent mixture, have been sound a solvent in some cases; as hath also spt. nitr. dulcis, the solution, or will of juniper. In violent sits of the disorder, the semicupium is good; and clysters with Venice turpentine, or spirit of turpentine, and opium. In nausea and vomiting, the saline draughts, with tinct. theb. and care should be taken to keep the body properly open with cathartics of the cooling saline kind, or rather with manna, ol. ricini, and inf. sennæ, or other mild purge. Emuss. com. decoct. rad. alth. or the like, may be used freely as common drink. (See also Calic.)

STORAX.

Pectoral, vulnerary gr. v. ad 3j.

STRAINS.

If internal, give spermaceti, ball traumat or Irish slate, with spruce beer. If outward, apply lin. sapon vinegar, lin. vol. or arquebusade. If there be inflammation, soment and poultice bleed and purge.

STRANGURY.

SYMPTOMS. A difficulty of making water, attended with heat or scalding; often proceeds from blisters.

TREATMENT. Give emulf. com. or mucilage gum arabic freely. If these fail, boluses of camphire and opium. See also Urine.

STYPTICS

STYPTICS internal

Tinct. styptic 3 j. ad 3ss. Tinck. rosar. 3j. ad

zij, Elix. vitr. acid gut. x. ad xx.

External, alum, colcothar, aq. vitriol. cærul. acc. faturni, vitriol virid. flour, cobweb, dry lint, fur of hat, agaric, ball. traumatic.

SUGAR of Lead.

Externally cooling, aftringent.

SULPHUR Flowers.

Pectoral, alterative, aperient 3j. ad zij.

SULPHUR precipitated.

Ditto.

SULPHUR of Antimony, precipitated.

Alterative gr. ½ ad ij.

SWINE-POX. See Pox.

SYRUP of Buckthorn. Cathartic zj. ad ziij.

SYRUP of Kermes.

Cordial 3sf. ad 3ij.

SYRUP of Squilts.

Pectoral, diuretic 3st. ad zij.

SYRUP of white Poppies.

Anodyne zj. ad zij.

TABES DORSALIS.

SYMPTOMS. A wasting of the whole body, attended

attended with fever, but no spitting nor cough, by which it is known from a *Phthisis*. A collection of matter in some part of the body is usually the cause, but particularly attends the inflamation.

mation of a schirrous gland.

TREATMENT. If it arises from schirrous liver, foap, and mild chalybeates. If the schirrus, or abscess be external, open or extirpate. The bark is often proper in these cases, as is also myrrh with nitre, or tart. vitriolat. The body must be kept properly open; but purgings should be checked. Gentle exercise, wholesome air, and proper mild diet, with the use of asses milk, and the like, are often of much service.

TAMARINDS.

Cooling, antiseptic, ad libit.

Ditto, also opening zij. ad zss.

TARTAR Emetic.

Gr. j. ad iv. Febrifuge, diaphoretic gr. 1 ad gr. j.

TARTAR vitriolated.

Aperient, febrifuge, &c. 3ss. ad 3ij.

TEETHING.

SYMPTOMS. Inflammation and swelling of the gums, and slavering. Fever, and sometimes convulsions.

TREATMENT.

TREATMENT. Cooling febrifuges in case of ever; the body should be kept gently open. If onvulsions arise, give tinct. suling and carminatives, or ol. anisi, with magnetia. If a rash apears, Gascoign's powder, or pulv. contray. c. with nitre, are usually given. If a cough, spernaceti, or other balsamic pectoral; leeches and listers may be applied if necessary; and if the isorder is violent, the gums should be lanced. Gentle anodynes are often useful.

TENESMUS.

Symptoms. Frequent, or even continual nelination to go to stool, without occasion, or eing able to avoid any. Arising from the abration of mucus, &c. from the rectum and anus.

TREATMENT. Clysters, with mutton broth nd opium; or with starch, oil, and elect. e cord. or with spermaceti, suet, or the like. Bals. ocat. or spermaceti with tinct. theb. may also be iven inwardly.

TESTICLES inflamed.

TREATMENT. Bleed, purge, and even vomit, f necessary, in order to cause revulsion. Warm mollient somentations and poultices are usually ecommended; but Dr. Simmons, in his ingenious ractical treatise on the Genorrhæa, recommends old applications to the part, as being much nore essicacious. Febrisuges and gentle diahoretics, with an anodyne at night, are good.

If matter forms, open and let it out. The testii cles thould be suspended, and the patient kept as much as possible in an horizontal posture.

TETANOS.

of the muscles of the neck, and trunk of the body, drawing it backwards or forwards in a curve with convulsions.

opium, keeping the body open with ol. ricini, on other proper purge. The warm bath, emollient fomentations and fpirituous embrocations to the parts. Sal c. c. and the nervous medicines are good; and clysters with turpentine and opium. After the disorder, give bark, or other restoratives.

TETTERS.

TREATMENT. They may be touched with ink, a weak folution of green or white vitriol, facen. faturn. fal. tartari, or fublimate. If these prove too violent, foftening and cooling liniments, or cream. Ung. cærul. nit. spt. vin. camph. or liniments with white precipitate are good. Sometimes alteratives of the mercurial and antimonial kind, or cooling febrifuges and purges are necessary.

Pimples, Ringworms, Shingles, Grubs, Freckles, and other like cutaneous complaints, also give

way to some of the above methods.

THROAT, fore.
TREATMENT. A common inflammatory fore throat.

roat, proceeding from cold, &c. gives way to eeding, cooling purges, gentle sudorifics, acidated gargles, and lin. vol. applied outwardly. necessary, blisters may be applied behind the rs. If there are ulcers, bleeding and purging generally improper; warm, but mild and antitic diaphoretics, as rad. serp. virg. conf. mocrat. cortex, sal. c. c. contrayerva, &c. with tergent gargles, composed of tinct. myrrh, ney, &c. or the tinct. rosar. Blisters if necesy; and if much sever, proper cooling sebrifus. See also Angina and Quincy.

THRUSH.

SYMPTOMS. Little ulcers in the mouth and oat, of a white or yellow colour. Frequent children.

TREATMENT. Keep the body gently open th magnesia or rhubarb; give the saline mixe, with pulv. e ch. or contrayerva c. and ch the ulcers now and then with syr. moror. borax, or with alum, bol. arm. and honey, t. rosar. &c. See also Fever Apthose, Fever liary, Angina Maligna, and Venereal Disease.

TIN Filings.

Vermifuge zff. ad zij.

TIN pulverized. Ditto.

TINCTURE of Antimony.

Deobstruent, diuretic, diaphoretic gut. x. ad lx. etic 3ij. or more.

TINCTURE

TINCTURE aromatic. 3 j. ad 3 ij.

TINCTURE of Bark. Corroborant, febrifuge 3 j. ad 3ff.

TINCTURE of Bark, volatile. Ditte 3 ff. ad 3 j.

TINCTURE, bitter. Stomachic Zj. ad Zij.

TINCTURE of Cantharides. Diuretic, corroborant gut. x. ad xxx. or more

Nervous 3 ff. ad 3 ij.

TINCTURE of Ginnamon. Astringent, cordial 3 j. ad 3 ij.

TINCTURE Foetid. - of Saot.

Antispasmodic, &c. gut. xx. ad 3 j.

TINCTURE of Guaiacum, volatile. Antirheumatic, &c. gut. xx. ad 3 j. or more.

TINCTURE of black Hellebore. Deobstruent, aperient 3 sf. ad 3 ij.

TINCTURE of Jalap. Cathartic 3 ij. ad 5 vj.

TINCTURE of Japan Earth. Astringent 3 j. ad 3 ij.

TINCTURE

TINCTURE of Myrrh. Deobstruent, antiseptic gut. xv. ad 3 j.

TINCTURE of Rhubarb. Opening 3 ff. ad 3 ij. Stomachic, &c. 3 j. ad Ziij.

TINCTURA Sacra.

Opening zj. ad zij. Deobstruent Zij. ad zss.

TINCTURE Saturnine.

Aftringent, antiphthifical gut. xx. ad xxx:

TINCTURE of Sena. Opening and carminative zj. ad ziij.

TINCTURE of Snakeroot. Alexipharmic, Comachic, &c. 3st. ad 3 ij.

TINCTURE of Flowers of Steel. Deobstruent, corroborant,&c. gut. x. ad 3 ff.

TINCTURE of Steel with fpt. of Salt. Ditto, gut. v. ad xx.

TINCTURE, Styptic.

3 I. ad 3 j.
TINCTURE Thebaic.

Anodyne, &c. gut. v. ad 3 ff.

TINCTURE of Valerian. Nervous 3 ff. ad 3 i.

TINCTURE of Valerian, volatile. Gut, xx. ad lx.

TONICS:

TONICS.

Bark $\ni j$. ad $\Im j$. see also its preparations. Elix. vitriol gut. v. ad xx. Vin. chalyb. \Im ss. \Im see also Corroborants.

TOOTH-ACH.

TREATMENT. If hollow or decayed, apply balf. traumatic, ol. caryoph. origan. vel guaiaci, upon cotton to the part; or pills with camphire and opium, or rad. pyrethri. Some burn the nerve with ol. vitriol, aqua fortis, or an hot iron; or touch the Antithelix with the latter. If there be inflammation, bleeding, purging, and blifters behind the ears. If it proceeds from Cold or Rheumatifm, treat it as directed for those complaints.

TREACLE, Venice.

Alexipharmic, sudorific, &c. 3 ff. ad 3j.

TREMORS.

TREATMENT as in Nerveus Complaints and Pally.

TROC	CHES pe	Etoral, Teart-burn,	} ad l	ibitum.
TRO	OCHES	of Japan 1 of Squills.	Earth.	
-	`	of Sulphur		The state of the s
	-	of Vipers. of Nitre.		en e

Virtues as the respective simples, only in double or treble doses.

TUMOURS

TUMOURS Encysted.

Try to disperse them with mercurial unction or plaister, or friction; or if these fail, hemlock fomentations and poultices, and give ext. cicut. or mercurials, with sarsaparilla, &c. and occasional purges. If you cannot disperse them, endeavour at suppuration; if that fails, extirpation, where it can be done, is the only cure.

TURPENTINES.

Diurctic, detergent, ftrengthening Dj. ad Dj.

TUTTY.

Externally in collyriums for the eyes; cooling, aftringent.

TYMPANY.

A collection of wind or air in the intestines, particularly the colon, by which they become violently distended. It sometimes arises from checking a diarrhoea, or from the imprudent use of bark in intermittents.

REATMENT. If fever and full pulfe, bleed; keep the body open with gentle laxatives, joined with aromatics; but refrain from strong purges, and from carminatives not joined with proper aperients. The belly should be frequently rubbed, and swathed with slannel. Exercise is good; gentle corroborants, as flor. chamæm. with small doses of chalybeates joined with aromatics, are also very proper.

VALERIAN.

VALERIAN.

Nervous, antispasmodic, &c. Hs. ad 3 st. bis die.

VARIX, or Enlargement of a Vein.
TREATMENT. Keep it compressed with proper bandages, and if necessary, bleed and evacuate. If these fail, treat it as in the operation for Aneurism.

VENEREAL Disease.

Symptoms. A few days after an impure coition, a titillation about the glans penis; heat and pricking pain in making water; then a discharge of matter from the urethra, at first whitish, and in small quantity; afterwards yellow or green, or even reddish, and more copious. If it be suffered to proceed, inflammation and excoriation follow; painful erection or chordee. Sometimes phymosis (or an inability of drawing the ikin back over the glans) paraphymofis (or an inability of drawing the skin forward so as to cover the glans) swelling of the testicles, chancres, buboes, scabby eruptions, nocturnal pains, nodes, carious bones, ulcers in the throat and nose, &c.

TREATMENT. In the beginning, bleed, if nothing forbids; and give a cooling purge, a small dose of calomel, merc. calcinat. or other proper mercurial, every night; and nitre, with gum arabic and some gentle lenitive, several times in the day; repeating the purge occasionally. If these are not judged sufficient, a small quantity of

ung. cærul. fort. may be rubbed into the thigh or leg every night. The parts should be kept clean by washing them with warm milk and water. This course should be persisted in till the virulence is subdued, afterwards a few cooling purges; and balf. capiv. with aftringent, or other injections, if gleet or ulcer remain, will usually be sufficient.

Buboes are generally advised to be dispersed with ung. cærul. fort. but if they are advanced, they should be brought forward by proper suppuratives, opened and digested thoroughly off, which is usually attended with the most falutary effects. Abscesses in perinæo, should, if possible,

be difperfed.

Chancres may be anointed with ung. cærul. fort. after washing them with aq. phagedænic. For the Hernia Humoralis see Testicles, see also

Phymosis.

When the diforder is a confirmed Lues, we should either salivate, or at least keep the blood for some time highly charged with mercury; for either of which purpofes calomel, or merc. calcinat. &c. should be given inwardly; and 3 f. or 3 j. of ung. cærul. fort. rubbed every night into the thighs. In fhort, the whole dependence must be on mercury for destroying the virulency, other remedies being only had recourse to occasionally, according to the symptoms that arise. Thus, inflam natory symptoms are to be removed by bleeding, purging, or other proper medicines; K 2 ulcers,

red in such cases; but the disease which is the cause of these should be principally attended to, and when that is subdued by a proper mercurial course, the other symptoms will either disappear, or may be removed by requisite methods. A Gleet, for example, will sometimes remain after the disorder is cured; and must be treated with corroborants, and, if necessary, astringent injections; the cold bath, and a strengthening diet.

VERDIGRIS.

Externally detergent, good against fungous sless.

VERTIGO.

SYMPTOMS. Objects, though at rest, seem to turn round; dimness of sight, and sear of falling.

TREATMENT. Bleed in the jugular, and cup in the back part of the head; blifters kept open; a vomit and laxatives; then the nervous medicines, as pil. gum. caftor, fal c. c. &c. and chalybeate waters.

VINEGAR.

Cooling, antalkaline, antiseptic zj. ad zsss. VINEGAR distilled. Ditto.

VINEGAR Lithrage.
Cooling, aftringent, deobstruent gut. x. ad 3j.

VINEGAR of Squills.

Pectoral, diuretic, deobstruent gut. x. ad zj.

VIPERS.

VIPERS.

Nutritive, corroborant, as food or in broth.

VIPERS Fat.

Externally for bites of vipers, also ophthalmic.

VITRIOL, Blue.

Escarotic externally. Also emetic off. ad oi. Corroborant gr. 4 ad gr. j.

VITRIOL, Green.

As falt of steel.

VITRIOL, White.

Emetic Dj. Astringent gr. iij. ad Off. ernally cooling, aftringent.

ULCERS.

TREATMENT. Drefs tnem with proper digeftives; and if they are fungous, apply an escharotic: f foul, mix red precipitate with the digeftive, or use the mel Ægyptiacum, or tinct. myrrhæ; or elfe wash with aq. phagedæn. or the like. Dry lint alone, in some cases, is good. callous, cut or eat away the callosity. If the bone be carious, either scrape it, or make small perfoations to promote exfoliation; or dress with int, either dry, or wetted with tinct. myrrhæ. If humours abound, give proper alteratives and purges; and if putridity appears, the bark and elixir of vitriol.

VOMICA, or Abscess in the Lungs. SYMPTOMS. It is sometimes a consequence of pleurify;

pleurify; cough, difficult breathing, even after the pain which preceded it goes off; flight shiverings, followed by heat; a quick weak pulse; hectic fever; wasting of the body; the patient cannot lie down but on the side assected.

TREATMENT. Gentle medicines of the turpentine kind, balfamics and diuretics, steams of hot water, with balf. Tolu, benzoin, &c. received into the lungs; occasional laxatives, bleeding if inflammation, soft balfamic food, and gentle exercise.

VOMITING.

TREATMENT. If it ariseth from foulness of stomach (as bile, acidity, &c.) a vomit will be useful; and afterwards, if necessary, a rhubarb or other gentle purge; and stomachies, magnefia, opiates, &c. according to circumstances. If from spasns, the saline draughts, and proper opiates; medicines of the nervous class, joined with laxatives; and if necessary, a blifter to the back. If from inflammation, bleeding, cooling diluents and febrifuges, as the faline draughts with proper laxatives; and anodynes at discretion. If from weaknoss of the stomach, bark, bitters, or other corroborants; premifing a vomit if judged necessary. Chalybeate waters are also proper. If the stomach be fore, as it is termed, the balf. traumat: spermaceti, or other balfamics, are good. In vonitings, the best form of medicine is pills.

Vomiting.

Vomitings are often symptomatic, as in fits of the gravel, pregnancy, &c. the original disorder nutt, in those cases, claim the chief attention.

VOMITING of Blood. See Blood.

URINE bloody.

The treatment may be similar to that prescri-

URINE, profuse Discharge of, or Diabetes.

TREATMENT. As this complaint arises from relaxed state of the sphincter of the bladder, it nust be cured by proper corroborant remedies, as ark, tinct. rosar. tinct. stypt. elix. vitriol, alum, itters, and the like. Tinct. canthar. has been ound essicacious, as hath also vitriol cærul. in oses of half a grain, twice in a day. Exercise, nd light, but strengthening diet, are also proper.

URINE, Suppression of.

Called Iscuria, if total; Strangury, if but in art; and Distury, if the latter be attended with eat; and are usually occasioned by inflammation r spasses on the neck of the bladder, in consequence of blisters, or some other cause.

TREATMENT. Medicines with ol. amygd. and lenty of gum arabic, with opiates or camphire, necessary. The body should be kept opening the gentle laxatives, and modlient clysters; if eccssary, bleed, some at the helly, and advise the arm bath. If caruncles in the urethra are the:

caule,

cause, bougies, introduced with proper caution, are serviceable.

UTERUS inflamed, &c. See Womb.

WARTS.

Touch them with some caustic or escarotic daily, paring off first the decayed part; or cut them off with scissars, and touch the roots with blue vitriol, or the like. If it can conveniently be done, tie round the bottom a fine hair, or silk thread, and it will die away. The juice of celadine, or spurge rubbed over them, are esteemed good.

WATER Horfe-radifi. Diuretic, deobstruent zij. ad & st.

WATER, Lime, simple.

Alterative, antiscorbutic, antacid ziv. or more

WATER Phagedænic.
To wash foul ulcers.

WATER Sapphirine.

Detergent, aftringent.

WATERS simple, distilled.

The virtues of their respective plants, &c. Zij.

WATERS, Spirituous distilled.

The virtues of their respective ingredients gif. ad zj. Also cordial.

WATER

WATER fulphurated. Pectoral, cooling, &c. 3sf. ad 3 ij.

WATER Tar.

Alterative, deobstruent, corroborant iv. ad

WATER, Vitriolic, eamphorated. Ophthalmic.

WATER, Vitriol blue. Externally styptic.

WAX.

Balsamic, antidysenteric; externally healing.

WHEY, Alum.

Aftringent 3 iv. bis die. Externally aftringent, cooling.

WHEY, Scorbutic.

I iv. ad #ff. bis die.

WHITES. See Fluor albus.

WHITLOW.

TREATMENT similar to that directed for Boils and Abscesses.

WINE aloetic, alkaline.
Deobstruent, attenuant 3 j. ad 3 iij.

WINE antimonial.

Emetic 3 ij a 3 ss. Diaphoretic, diuretic, ebrifuge gut. x. ao ix.

WINE

WINE bitter.
Stomachic 3j. ad 3iij. bis die.

WINE Ipecacuanha.

Emetic zff. ad zij.

WINE of Millepedes.

Diuretic, attenuant, decbstruent zj. ad ziij.

WINE, Saffron.

Cordial zij. ad zis.

WINE, Steel.

Deobstruent, corroborant 3 sf. ad 3 iij. bis die.

WINE, Viper.

Strengthening zij. ad 3j.

WOMB inflamed.

SYMPTOMS. Pulsation, pain, heat, and tumour in the part: fever, costiveness, pain on going to

stool, and difficulty of making water.

TREATMENT. Bleeding, laxatives, emollient elysters, cooling febrifuges, diluting liquors, somentations and poultices, &c. as in other inflammations. (See Stomach, Bladder, Kidnies, &c.)

If a Cancer fucceeds, it must be palliated with gentle anodynes, occasional bleeding and laxatives; and proper detergent, or other injections.

WORMS.

SYMPTOMS. Itching of the note and anus; hun-

ger, with ravenous appetite; nausea; and gnawing pain in the stomach or intestines; social breath; grinding of the teeth, and moaning in sleep; intermitting weak pulse, and hectic sever; faintings, and sometimes convulsive sits; palenes; wasting of the slesh. But there are several kinds of worms. Itching about the anus, with tenesmus, are signs of ascarides, or small thread worms. Sudden gripings about the navel, denote the common round worms. Gnawing pains in the stomach, and voraciousness, the maw worms. And a weight in the belly, like the rolling of a

ball, the teneâ, or tape worm.

TREATMENT. Tin filings, with rhubarb, or other proper aperient; sæm. santon. calomel, æthiops, or other mecurials, every night; with a purge now and then. Indian pink-root off. in powder, or corallin. The infusion of bears-foot leaves is esteemed good; cabbage-tree bark has also been recommended. Aloes, wormwood, rue, savine, especially in clysters; and for the ascarides, oil of olives, or the sumes of tobacco thrown up the rectum; for the tape worm, the male fern root zj. every hour for three times, and presently after a strong drastic mercurial purge. See Dr. Simmons's ingenious pamphlet on the subject. Corroborants are proper after a course of anthelmintic medicines.

WORMWOOD.

In clysters, vermifuge,

WORMWOOD

WORMWOOD in Infusion.

Stomachic, externally in fomentations discutient: WORM Seed.

Vermifuge gr. v. ad Dj. or more in infusion. WOUNDS.

TREATMENT. Remove extraneous subfrances; and, if the wound be simple, close the lips with slips of sticking plaisser, or the stitch if necessary; and apply lint, either dry, or in case of bleeding, wetted with ball, traumatic, &c. If necessary, bleed and purge; or give cordial, &c. according to circumstances. Dress with some mild digestive, or defensative; and if sungous sless appears.

use escarotics, afterwards cicatrize.

If there be loss of flesh, defend the part from the air with proper, mild, vulnerary dreffings, while nature supplies the loss. In other respects (the future excepted) proceed as above. If an artery or vein be wounded or divided, apply Myptics, ligatures, &c. according to circumstanrces. If a nerve be divided in part (as violent pain, convultions, and dilirium will indicate) divide it intirely; as must also be the case with a tendon or artery in like circumstances if there be fever, bleed, purge, and give cooling febrifuges; and if inflammation, apply emolliant poultices and for entations. If the patient is low, and the wound does not go on kindly, bark, or other proper cordial and threngthening medicines, with fuitable diet, thould be prescribed. TABULA

TABULA oftendens, quâ proportione fimplica medicamenta purgantia, narcotica, & mercurialia, in compositis, quibus occurrunt, continentur.

Pulvis e bolo compositus cum opio in gr. 45. Pulvis e scammonio compositus in gr. 7. Pulvis e senà compositus in gr. 21.

Pulvis e fuccino compofitus in gr. 40.

pecies e feordio cum
opio in gr. 45.

Pilulæ ex colocynthide
fimpliciores in zff.

ilulæ ex colocynthide cum aloe in 3ff.

Hulæ mercuialet in gr. 28.

continet Opii gr. 1. continet Scammonii gr. 4 continet Senæ gr. 8. Crystallorum tartari gr. 2. Scammoni gr. 2. continet Opii gr. I. continent Opii gr. 1. continent fere Scammonii off. Colocynthidis off. continent ferè Aloes gr. 8. Scammonii gr. 8. Colocynthidis gr. 4. continent Argenti vivi gr. 15.

Pilulæ

Pilulæ saponaceæ in Dst. continent ferè Opii gr. I. Pilulæ e styrace in gr. continent Opii gr. Takana 54. Electarium e scammonio continet in ziss. Scammonii gr. 15. continet Electarium e scordio in Opii gr. I. ziii. continet . Confectio paulina in gr. Opii gr. 1. 32. continet . . Mithridatium, five con-Opii gr. I. fectio Damocratis in 3ff. Philonium Londinense in continet gr. 36. Opii gr. I. Theriaca Andromachi in continet Opii gr. 1.

In Emplastro ex ammoniaco cum mercurio Ar-

gentum vivum est totius pars ferè quinta.

In Emplastro communi cum mercurio Argentum vivum est totius pars serè quinta.

In Unguento cæruleo fortiori Argentum

vivum est totius pars ferè tertia.

In Unguento cæruleo mitiori Argentum

vivum est totius pars ferè quinta.

In Cerato mercuriali Argentum vivum est totius pars ferè quinta.











